



OPEN ACCESS

APPROVED BY

Frontiers Editorial Office, Frontiers Media SA, Switzerland

*CORRESPONDENCE Ali Mekki Missaoui, ☑ cssamm@uga.edu

RECEIVED 24 October 2025 REVISED 04 November 2025 ACCEPTED 04 November 2025 PUBLISHED 12 November 2025

CITATION

Makaju SO, Chhetri HB, Abeyratne CR, Pavicic M, Poudel H, Irfan JA, Giabardo A, Devos KM, Jacobson D and Missaoui AM (2025) Correction: Drought adaptation index (DAI) based on BLUP as a selection approach for drought-resilient switchgrass germplasm. *Front. Genet.* 16:1731825. doi: 10.3389/fgene.2025.1731825

COPYRIGHT

© 2025 Makaju, Chhetri, Abeyratne, Pavicic, Poudel, Irfan, Giabardo, Devos, Jacobson and Missaoui. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Correction: Drought adaptation index (DAI) based on BLUP as a selection approach for drought-resilient switchgrass germplasm

Shiva Om Makaju^{1,2,3}, Hari Bahadur Chhetri⁴, Chanaka Roshan Abeyratne⁴, Mirko Pavicic⁴, Hari Poudel⁵, Jazib Ali Irfan^{2,3}, Anita Giabardo², Katrien M. Devos^{2,3,6}, Daniel Jacobson⁴ and Ali Mekki Missaoui^{1,2,3}*

¹Center for Applied Genetic Technologies, University of Georgia, Athens, GA, United States, ²Institute of Plant Breeding, Genetics and Genomics, University of Georgia, Athens, GA, United States, ³Department of Crop and Soil Sciences, University of Georgia, Athens, GA, United States, ⁴Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN, United States, ⁵Agriculture and Agri-Food Canada, Lethbridge Research and Development Centre, Lethbridge, AB, Canada, ⁶Department of Plant Biology, University of Georgia, Athens, GA, United States

KEYWORDS

switchgrass, bioenergy, drought adaptation, yield stability, BLUP, drought adaptation index, stress tolerance indices, biomass

A Correction on

Drought adaptation index (DAI) based on BLUP as a selection approach for drought-resilient switchgrass germplasm

by Makaju SO, Chhetri HB, Abeyratne CR, Pavicic M, Poudel H, Irfan JA, Giabardo A, Devos KM, Jacobson D and Missaoui AM (2025). Front. Genet. 16:1626083. doi: 10.3389/fgene. 2025.1626083

The following supplemental materials were omitted.

SUPPLEMENTARY FIGURE 1 | Spatial plots for each year from 2019 through 2022. CV and UC represent drought and control fields, respectively.

SUPPLEMENTARY FIGURE 2 | Histogram of raw dry biomass yield (g plant $^{-1}$) before the removal of yield data rows below 50 g plant $^{-1}$. CV and UC represent drought and control fields, respectively.

SUPPLEMENTARY FIGURE 3 | Histogram of raw dry biomass yield (g plant⁻¹) after the removal of yield data rows below 50 g plant⁻¹. CV and UC represent drought and control fields, respectively.

SUPPLEMENTARY FIGURE 4 | QQ plot of data after outlier removal.

SUPPLEMENTARY FIGURE 5 | Distribution of BLUP predicted values.

SUPPLEMENTARY FIGURE 6 | Biplot of genotype mean performance (yield as BLUP-predicted value) versus second interaction principal component (IPCA2) scores.

The file(s) have now been published.

The original article has been updated.

Makaju et al. 10.3389/fgene.2025.1731825

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.