



## OPEN ACCESS

## EDITED BY

Antonio Sarria-Santamera,  
Nazarbayev University, Kazakhstan

## REVIEWED BY

Jun-Yu Ma,  
Guangdong Second Provincial General  
Hospital, China  
Sharapat Moynbayeva,  
Kazakhstan School of Public Health,  
Kazakhstan

## \*CORRESPONDENCE

Leila Stabayeva

✉ stabaeva@qmu.kz

Olzhas Zhamantayev

✉ zhamantaev@qmu.kz

RECEIVED 25 April 2025

ACCEPTED 04 September 2025

PUBLISHED 23 September 2025

## CITATION

Podilyakina Y, Stabayeva L, Kulov D,  
Kamyshanskiy Y, Amirbekova Z, Stundžienė R  
and Zhamantayev O (2025) Risk factors for  
first-trimester spontaneous abortion and the  
role of preconception care.  
Front. Glob. Women's Health 6:1615983.  
doi: 10.3389/fgwh.2025.1615983

## COPYRIGHT

© 2025 Podilyakina, Stabayeva, Kulov,  
Kamyshanskiy, Amirbekova, Stundžienė and  
Zhamantayev. This is an open-access article  
distributed under the terms of the [Creative  
Commons Attribution License \(CC BY\)](#). The  
use, distribution or reproduction in other  
forums is permitted, provided the original  
author(s) and the copyright owner(s) are  
credited and that the original publication in  
this journal is cited, in accordance with  
accepted academic practice. No use,  
distribution or reproduction is permitted  
which does not comply with these terms.

# Risk factors for first-trimester spontaneous abortion and the role of preconception care

Yuliya Podilyakina<sup>1</sup>, Leila Stabayeva<sup>2\*</sup>, Dusentay Kulov<sup>3</sup>,  
Yevgeniy Kamyshanskiy<sup>4</sup>, Zhanna Amirbekova<sup>5</sup>,  
Rasa Stundžienė<sup>6</sup> and Olzhas Zhamantayev<sup>3\*</sup>

<sup>1</sup>School of Residency and Professional Development, Karaganda Medical University, Karaganda, Kazakhstan, <sup>2</sup>Department of Morphology, Karaganda Medical University, Karaganda, Kazakhstan, <sup>3</sup>School of Public Health, Karaganda Medical University, Karaganda, Kazakhstan, <sup>4</sup>Clinic of the Karaganda Medical University, Karaganda, Kazakhstan, <sup>5</sup>Department of Obstetrics and Gynecology, Karaganda Medical University, Karaganda, Kazakhstan, <sup>6</sup>Faculty of Medicine, Institute of Health Sciences, Department of Nursing, Vilnius University, Vilnius, Lithuania

**Background:** Spontaneous abortion in the first trimester is a common adverse pregnancy outcome with significant implications for maternal health and public health practice. The description of associations with modifiable factors, including preconception care, can aid in planning strategies to improve pregnancy outcomes.

**Methods:** A retrospective analysis was conducted using data from 1,526 women, divided into two groups based on pregnancy outcomes: spontaneous abortion in the first trimester and live births. Binary and multivariate logistic regression analyses were performed to identify associations between factors (including preconception care) and the risk of spontaneous abortion in the first trimester.

**Results:** Age >35 years [OR] = 2.02, 95% [CI] = 1.49–2.75, obesity [OR] = 1.81, 95% [CI] = 1.12–2.91, and a history of spontaneous abortion [OR] = 1.57, 95% [CI] = 1.01–2.43 were associated with higher odds of spontaneous abortion in the first trimester, whereas preconception care was associated with lower odds of spontaneous abortion in the first trimester [OR] = 0.58, 95% [CI] = 0.45–0.75.

**Conclusion:** The findings may help clinicians stratify pregnant women who require additional monitoring and pre-pregnancy interventions. From a public health perspective, integrating preconception care into routine health services can enhance maternal and neonatal outcomes, reduce healthcare costs, and improve health equity by targeting vulnerable populations. However, the results should be interpreted as associations, and prospective studies are needed to assess the potential effects of preconception care on spontaneous abortion in the first trimester.

## KEYWORDS

women's health, reproductive health, preconception care, spontaneous abortion, miscarriage, maternal health

# 1 Introduction

Spontaneous abortion (SA), classified under ICD-10 code O03 and commonly referred to as miscarriage, is a prevalent complication in early pregnancy, with incidence rates varying depending on population characteristics and study methodologies (1, 2). Research indicates that approximately 10% to 20% of clinically recognized pregnancies worldwide result in SA, with higher rates observed when early pregnancy losses are included (3, 4). The majority of these losses occur in the first trimester (5, 6). In Kazakhstan, a study involving 237 participants reported that 25.6% of respondents had experienced SA (7). A large-scale national study analyzing 207,317 records of women across all regions of Kazakhstan estimated the prevalence of spontaneous pregnancy loss before 22 weeks of gestation to be 8.7%, with a 20% decline observed over the 2014–2019 period (8).

Several factors have been associated with an increased risk of SA. Advanced maternal age is a significant risk factor, as chromosomal abnormalities in embryos are more common in older women (9, 10). A cross-sectional study in Astana, Kazakhstan (2015–2017), analyzing 67,759 inpatient records, found advancing maternal age significantly increased the risk of miscarriage, with women aged 30–39 and 40+ years having 54% and 272% higher risks, respectively compared to women aged 19–29 years (11). Moreover, maternal obesity has been linked to higher miscarriage rates, potentially due to hormonal imbalances and metabolic disturbances, with risks ranging from 1.25 to 2.25 times higher compared to normal-weight women (12–14). Another important factor in early pregnancy loss is the history of SA and pre-pregnancy health. Studies have shown that a history of SA significantly increases the risk of subsequent losses, with women who have experienced one or more miscarriages being at higher risk for recurrence (15, 16). Inadequate pre-pregnancy health, including suboptimal management of chronic conditions, poor nutritional status, and lack of preconception care, has been linked to adverse pregnancy outcomes (17). Lifestyle factors prior pregnancy, including smoking and alcohol consumption, have also been implicated in elevating miscarriage risk (18, 19).

Building on these findings, it becomes clear that preconception may play an important role in addressing the modifiable risk factors associated with early pregnancy loss (20). Preconception care, as defined by the World Health Organization, includes a set of biomedical, behavioral, and social interventions provided to women and couples before conception to improve maternal and fetal outcomes (21). It aims to identify and mitigate risk factors such as obesity, unmanaged chronic conditions, and inadequate nutritional status while promoting healthier lifestyle choices (22). The benefits of preconception care are well-documented, with evidence indicating that it can help reducing the incidence of adverse pregnancy outcomes, including SA, preterm birth, and congenital anomalies (23). Targeted preconception interventions, such as weight management, smoking cessation, and folic acid supplementation, have been shown to improve pregnancy outcomes and reduce

healthcare costs by preventing complications before conception (24).

This study aimed to investigate the socio-demographic and clinical risk factors associated with SA in the first trimester and evaluate the protective effects of preconception care among women in Kazakhstan, largest country in Central Asia. It adds a global health perspective and highlights the importance of addressing disparities in maternal health outcomes, an essential aspect of public health research. While previous studies have extensively documented risk factors for spontaneous abortion, there is limited research on the role of preconception care in mitigating these risks, particularly in low- and middle-income countries (LMICs), and some finding can be extrapolated to all Central Asian countries.

# 2 Materials and methods

## 2.1 Study design

This retrospective study included women who received health services for SA in the first trimester or during childbirth at the State Clinical Hospital, a large conglomerate that includes two perinatal centers of the first and second levels in Karaganda, Kazakhstan, from January 1, 2018, to January 1, 2023. Women from across the country, including those with complicated medical histories, are admitted to these perinatal centers, as they provide the most advanced medical care. The first-level perinatal center offers basic obstetric and neonatal care for low-risk pregnancies and routine deliveries, while the second-level perinatal center provides specialized care for moderate and high-risk pregnancies and newborns requiring additional medical attention.

The use of clinical data from women and morphological data of SA tissue for research purposes was approved by the Local Commission on Bioethics of Karaganda Medical University. Written informed consent was obtained from all pregnant women for the use of their clinical and morphological data in the study.

Women's clinical and sociodemographic data were obtained from the Integrated Health Information System (IHIS) and reviewed for missing values. Patients with incomplete data were excluded from the final analysis (Supplementary Table S1). Women with ectopic pregnancies, multiple pregnancies, or antenatal or intrapartum fetal deaths were also excluded. The study block diagram is presented in Figure 1.

The study included data from 1,526 women, divided into two groups:

1. Favorable outcome ( $n = 1,236$ ): Pregnancy resulting in live birth.
2. SA in the first trimester ( $n = 290$ ): Pregnancy ending in spontaneous abortion.

Variables extracted from IHIS included women's age at the time of the study, ethnicity, place of residence, social status, education, marital status, body mass index (BMI), type 1 or type 2




FIGURE 1

Flowchart illustrating the recruitment process of study participants.

diabetes, arterial hypertension, hypothyroidism, deficiency anemia, allergies, health insurance status before pregnancy, parity (divided into two groups), and preconception care. Preconception care interventions included nutritional counseling, folic acid supplementation, management of chronic conditions (e.g., diabetes, hypertension), and lifestyle modifications (e.g., weight management). These services were provided to women who sought care at the State Clinical Hospital in Karaganda.

The primary outcome variable was the pregnancy outcome, classified into two groups: “First-trimester SA” and “Favorable

outcome.” A favorable outcome was defined as the birth of a live newborn. The diagnosis of first-trimester SA was made clinically, based on bleeding and apparent expulsion of the embryo or fetus (confirmed by histological examination), or through ultrasound.

The first trimester was chosen as the cut-off point because SA occurring before this period typically shares a common pathophysiological etiology, such as inflammation, luteal phase deficiency, or chromosomal and structural abnormalities, which differ from miscarriages occurring after the first trimester.

The date of birth of each study participant was used to calculate their age. Age was classified into three categories: 18–26 years, 27–35 years, and over 35 years. Minors were excluded from the study.

## 2.2 Definitions

**First-trimester SA:** Spontaneous abortion occurring before the 13th week of gestation (25, 26).

**Favorable outcome:** Defined in this study as a pregnancy resulting in live birth.

In this study, preconception care referred to a set of measures conducted six or more months before pregnancy and documented in patients' medical records. These included nutritional counseling (with recommendations for folic acid intake), screening and management of chronic diseases as indicated, medical-genetic counseling when necessary, testing for HIV infection, hepatitis B and C, and sexually transmitted infections, as well as recommendations for lifestyle modifications for both spouses (body weight control, physical exercise, and cessation of harmful habits). These medical services were provided in accordance with the recently updated standard of obstetric and gynecological care in the Republic of Kazakhstan (27).

**BMI before pregnancy:** Calculated as the registered weight (kg) divided by the square of measured height (m) and categorized into four groups. For Caucasians: underweight ( $<18.5 \text{ kg/m}^2$ ), normal weight ( $\geq 18.5$  and  $<25 \text{ kg/m}^2$ ), overweight ( $\geq 25$  and  $<30 \text{ kg/m}^2$ ), and obese ( $\geq 30 \text{ kg/m}^2$ ). For Asians: underweight ( $<18.5 \text{ kg/m}^2$ ), normal weight ( $\geq 18.5$  and  $<22.9 \text{ kg/m}^2$ ), overweight ( $\geq 23$  and  $<24.9 \text{ kg/m}^2$ ), and obese ( $\geq 25 \text{ kg/m}^2$ ) (28, 29).

**Diabetes mellitus** was diagnosed based on  $\text{HbA1c} \geq 6.5\%$  (48 mmol/mol) and/or fasting plasma glucose or 2 h post-oral glucose tolerance test glucose levels  $\geq 7.0 \text{ mmol/L}$ .

**Hypertension:** Defined as a systolic blood pressure  $\geq 140 \text{ mmHg}$  or a diastolic blood pressure  $\geq 90 \text{ mmHg}$  in two separate measurements taken before 20 weeks of gestation (30).

**Hypothyroidism** was identified based on a pre-pregnancy diagnosis with thyroid-stimulating hormone (TSH) levels  $\geq 2.5 \text{ mIU/L}$  or, in the first trimester,  $\text{TSH} > 4 \text{ mIU/L}$ .

**Anemia:** Defined as hemoglobin (Hb)  $< 110 \text{ g/L}$  and hematocrit  $< 33\%$ , or as a transferrin saturation (TSat)  $< 16\%$ , ferritin concentration  $< 30 \text{ } \mu\text{g/L}$ , or vitamin B12 concentration  $< 200 \text{ pg/ml}$  (31).

## 2.3 Statistical analysis

Statistical analysis was performed using IBM SPSS Statistics (version 25.0, IBM Corp., Armonk, NY, USA). Percentages and frequencies were calculated for categorical variables, while the normality of continuous variables was tested using the Kolmogorov–Smirnov test. The Mann–Whitney  $U$  test was used to examine group differences for continuous variables. For

categorical variables, comparisons were made using the chi-square test.

Associations between potential factors (including preconception care) and first-trimester spontaneous abortion were assessed using univariable and multivariable logistic regression. The outcome was coded as  $\text{SA} = 1$  (adverse outcome). Discrimination of the final adjusted multivariable model was evaluated using the area under the receiver operating characteristic curve (AUC), calculated based on the predicted probabilities of this model [standard error [SE] and 95% confidence interval [CI] are reported]. Model calibration was assessed using the Hosmer–Lemeshow test.

Variables with  $p < 0.10$  in univariable analysis were considered for inclusion in the multivariable model. A purposeful selection approach followed: variables that were non-significant and did not substantially influence the coefficients of key predictors were sequentially excluded. The final adjusted model included age over 35 years, obesity, history of spontaneous abortion, and preconception care. The AUC was calculated based on the predicted probabilities of this final model (outcome  $\text{SA} = 1$ ).

Multicollinearity was assessed using variance inflation factors (VIF), and all predictors had acceptable VIF values ( $< 2$ ). The linearity of continuous variables was confirmed through scatterplot analysis. Odds ratios (OR) with 95% confidence intervals (CI) were used to report the results of the model.

## 3 Results

### 3.1 Socio-demographic and clinical characteristics of the original sample

Table 1 presents the socio-demographic and clinical characteristics of the sample. More than half of the sample, 775 women (50.8%), were aged 27–35 years, one-third, 476 women (31.2%), were aged 18–26 years, and 275 women (18%) were over 35 years old. Regarding ethnic composition, Kazakhs accounted for 732 women (48%), Russians for 595 women (39%), and other nationalities for 199 women (13%). A majority, 1,172 women (76.8%), lived in urban areas, while 354 women (23.2%) lived in rural areas.

In terms of social status, 683 women (44.8%) were classified as “other,” 405 women (26.5%) were employed (including civil servants, state employees, employees of private organizations, and workers), 23 women (1.5%) were unemployed, and 411 women (26.9%) were housewives. At the time of the study, 217 women (14.2%) had secondary education, while 1,309 women (85.8%) had secondary specialized or higher education. Additionally, 1,267 women (83%) were married.

Among the women, 946 (62%) had a normal weight, 213 (14%) were underweight, 274 (18%) were overweight, and 93 (6%) suffered from first- or second-degree obesity. A history of diabetes mellitus type 1 or 2 was reported in 49 women (3.2%), arterial hypertension in 56 women (3.7%), hypothyroidism in 26

**TABLE 1** Socio-demographic and clinical characteristics of study participants.

| Variables                       | Socio-demographic characteristics | All participants (n = 1,526) |
|---------------------------------|-----------------------------------|------------------------------|
| Woman's age, years              | 18–26                             | 476 (31.2%)                  |
|                                 | 27–35                             | 775 (50.8%)                  |
|                                 | >35                               | 275 (18.0%)                  |
| Ethnicity                       | Kazakh                            | 732 (48.0%)                  |
|                                 | Russian                           | 595 (39.0%)                  |
|                                 | Others                            | 199 (13.0%)                  |
| Residency                       | Urban                             | 1,172 (76.8%)                |
|                                 | Rural                             | 354 (23.2%)                  |
| Social status                   | Employed, civil servant           | 87 (5.7%)                    |
|                                 | Employed, public sector worker    | 111 (7.3%)                   |
|                                 | Employed, private sector worker   | 59 (3.9%)                    |
|                                 | Employed, self-employed           | 27 (1.8%)                    |
|                                 | Employed, laborer                 | 9 (0.6%)                     |
|                                 | Employed, other                   | 112 (7.3%)                   |
|                                 | Convicted                         | 4 (0.3%)                     |
|                                 | Other                             | 683 (44.8%)                  |
|                                 | Unemployed                        | 23 (1.5%)                    |
|                                 | Housewife                         | 411 (26.9%)                  |
| Education                       | Secondary education               | 217 (14.2%)                  |
|                                 | Higher than secondary education   | 1,309 (85.8%)                |
| Marital status                  | Married                           | 1,267 (83.0%)                |
|                                 | Not married                       | 259 (17.0%)                  |
| <b>Clinical characteristics</b> |                                   |                              |
| BMI, kg/m <sup>2</sup>          | Underweight                       | 213 (14.0%)                  |
|                                 | Normal weight                     | 946 (62.0%)                  |
|                                 | Overweight                        | 274 (18.0%)                  |
|                                 | Obese                             | 93 (6.0%)                    |
| Type 1/ Type 2 diabetes         | Yes                               | 49 (3.2%)                    |
|                                 | No                                | 1,477 (96.8%)                |
| Arterial hypertension           | Yes                               | 56 (3.7%)                    |
|                                 | No                                | 1,470 (96.3%)                |
| Hypothyroidism                  | Yes                               | 26 (1.7%)                    |
|                                 | No                                | 1,500 (98.3%)                |
| Deficiency anemia               | Yes                               | 382 (25.0%)                  |
|                                 | No                                | 1,144 (75.0%)                |
| Allergies                       | Yes                               | 68 (4.5%)                    |
|                                 | No                                | 1,458 (95.5%)                |
| Pre-pregnancy health insurance  | Yes                               | 1,329 (87.1%)                |
|                                 | No                                | 197 (12.9%)                  |
| Parity                          | Primiparous                       | 919 (60.2%)                  |
|                                 | Multiparous                       | 607 (39.8%)                  |
| History of SA                   | Yes                               | 122 (8.0%)                   |
|                                 | No                                | 1,404 (92.0%)                |
| Preconception care              | Yes                               | 930 (60.9%)                  |
|                                 | No                                | 596 (39.1%)                  |

BMI, body mass index; SA, spontaneous abortion.

women (1.7%), and deficiency anemia in 382 women (25%). Additionally, 68 women (4.5%) suffered from allergies.

The majority, 1,329 women (87.1%), had medical insurance before pregnancy. This pregnancy was the first for 919 women (60.2%). A history of early pregnancy loss was reported in 122 women (8.0%). Moreover, 930 women (60.9%) had undergone preconception training.

### 3.2 Comparative groups of women with a favorable pregnancy outcome and first-trimester SA

The socio-demographic and clinical characteristics of women classified according to pregnancy outcome are presented in **Table 2**.

In the group with first-trimester SA, compared to women with a favorable pregnancy outcome, there was a higher proportion of women over 35 years of age (26.9% vs. 15.9%). Additionally, women with SA were more likely to live in urban areas (79.7% vs. 76.1%), have higher than secondary education (86.2% vs. 85.7%), and be unmarried (17.6% vs. 16.8%).

Clinically, women in the first-trimester SA group had a higher prevalence of obesity (9.3% vs. 5.3%), type 1 or 2 diabetes (5.2% vs. 2.8%), arterial hypertension (5.2% vs. 3.3%), and allergies (5.9% vs. 4.1%). Furthermore, a larger proportion of these women had a history of SA (11% vs. 7.3%). In contrast, relatively fewer women in the SA group underwent preconception care compared to those with a favorable pregnancy outcome (49.7% vs. 63.6%).

The results of the univariate analysis of risk factors for SA in the study sample are presented in **Table 3**.

Univariate logistic regression analysis revealed that women aged over 35 years [[OR] = 1.94, 95% [CI] = 1.44–2.62], obesity [[OR] = 1.82, 95% [CI] = 1.14–2.90], type 1 or type 2 diabetes [[OR] = 1.93, 95% [CI] = 1.04–3.59], and a history of miscarriage [[OR] = 1.58, 95% [CI] = 1.03–2.42] were associated with an increased risk of first-trimester SA. Preconception care, however, was associated with a reduced risk of first-trimester SA [[OR] = 0.57, 95% [CI] = 0.44–0.73].

**Figure 2; Table 3** summarize the results of the multivariate logistic regression analysis. Independent factors influencing first-trimester SA included the woman's age, BMI, history of SA, and preconception care ( $p < 0.05$ ). Women aged over 35 had an increased likelihood of first-trimester SA compared to those under 35 [[OR] = 2.02, 95% [CI] = 1.49–2.75]. Obese women were more likely to experience first-trimester SA than women with a normal BMI [[OR] = 1.81, 95% [CI] = 1.12–2.91]. The history of SA was also strongly associated with a higher risk of first-trimester SA [[OR] = 1.57, 95% [CI] = 1.01–2.43]. Women who received preconception care had a significantly reduced risk of first-trimester SA compared to those who did not [[OR] = 0.58, 95% [CI] = 0.45–0.75].

The final adjusted model included age over 35 years, obesity, history of spontaneous abortion, and preconception care. The discriminative ability of the model was moderate: AUC = 0.611 (SE 0.019; 95% CI 0.574–0.648;  $p < 0.001$  compared with 0.50). Model calibration was satisfactory: Hosmer–Lemeshow test  $\chi^2 = 2.04$ , df = 4,  $p = 0.73$ .

## 4 Discussion

For practical healthcare, a key conceptual issue is the variability of risk factors for spontaneous miscarriage, as



TABLE 2 Comparative groups of women with a favorable pregnancy outcome and first-trimester SA.

| Variables                         |                                 | Favorable outcome<br>( <i>n</i> = 1,236) | SA in the first trimester<br>( <i>n</i> = 290) | <i>p</i> -value |
|-----------------------------------|---------------------------------|------------------------------------------|------------------------------------------------|-----------------|
| Socio-demographic characteristics |                                 |                                          |                                                |                 |
| Age, years                        | 18–26                           | 399 (32.3%)                              | 77 (26.6%)                                     | <0.001*         |
|                                   | 27–35                           | 640 (51.8%)                              | 135 (46.6%)                                    |                 |
|                                   | >35                             | 197 (15.9%)                              | 78 (26.9%)                                     |                 |
| Ethnicity                         | Kazakh                          | 593 (48.0%)                              | 139 (47.9%)                                    | 0.725           |
|                                   | Russian                         | 478 (38.7%)                              | 117 (40.4%)                                    |                 |
|                                   | Other                           | 165 (13.3)                               | 34 (11.7%)                                     |                 |
| Residency                         | Urban                           | 941 (76.1%)                              | 231 (79.7%)                                    | 0.201           |
|                                   | Rural                           | 295 (23.9%)                              | 59 (20.3%)                                     |                 |
| Social status                     | Employed, civil servant         | 70 (5.7%)                                | 17 (5.9%)                                      | 0.724           |
|                                   | Employed, public sector worker  | 92 (7.4%)                                | 19 (6.6%)                                      |                 |
|                                   | Employed, private sector worker | 48 (3.9%)                                | 11 (3.8%)                                      |                 |
|                                   | Employed, self-employed         | 21 (1.7%)                                | 6 (2.1%)                                       |                 |
|                                   | Employed, laborer               | 8 (0.6%)                                 | 1 (0.3%)                                       |                 |
|                                   | Employed, other                 | 91 (7.4%)                                | 21 (7.2%)                                      |                 |
|                                   | Convicted                       | 4 (0.3%)                                 | 0 (0%)                                         |                 |
|                                   | Other                           | 549 (44.4%)                              | 134 (46.2%)                                    |                 |
|                                   | Unemployed                      | 19 (1.5%)                                | 4 (1.4%)                                       |                 |
|                                   | Housewife                       | 334 (27.0%)                              | 77 (26.6%)                                     |                 |
| Education                         | Secondary education             | 177 (14.3%)                              | 40 (13.8%)                                     | 0.818           |
|                                   | Higher than secondary education | 1,059 (85.7%)                            | 250 (86.2%)                                    |                 |
| Marital status                    | Married                         | 1,028 (83.2%)                            | 239 (82.4%)                                    | 0.758           |
|                                   | Not married                     | 208 (16.8%)                              | 51 (17.6%)                                     |                 |
| Clinical characteristics          |                                 |                                          |                                                |                 |
| BMI, kg/m <sup>2</sup>            | Underweight                     | 168 (13.6%)                              | 45 (15.5%)                                     | 0.053*          |
|                                   | Normal weight                   | 777 (62.9%)                              | 169 (58.2%)                                    |                 |
|                                   | Overweight                      | 225 (18.2%)                              | 49 (17.0%)                                     |                 |
|                                   | Obesity                         | 66 (5.3%)                                | 27 (9.3%)                                      |                 |
| Type 1/Type 2 diabetes            | Yes                             | 34 (2.8%)                                | 15 (5.2%)                                      | 0.036*          |
|                                   | Not                             | 1,202 (97.2%)                            | 275 (94.8%)                                    |                 |
| Arterial hypertension             | Yes                             | 41 (3.3%)                                | 15 (5.2%)                                      | 0.131           |
|                                   | No                              | 1,195 (96.7%)                            | 275 (94.8%)                                    |                 |
| Hypothyroidism                    | Yes                             | 21 (1.7%)                                | 5 (1.7%)                                       | 0.977           |
|                                   | No                              | 1,215 (98.3%)                            | 285 (98.3%)                                    |                 |
| Deficiency anemia                 | Yes                             | 310 (25.1%)                              | 72 (24.8%)                                     | 0.929           |
|                                   | No                              | 926 (74.9%)                              | 218 (75.2%)                                    |                 |
| Allergies                         | Yes                             | 51 (4.1%)                                | 17 (5.9%)                                      | 0.198           |
|                                   | No                              | 1,185 (95.9%)                            | 273 (94.1)                                     |                 |
| Pre-pregnancy health insurance    | Yes                             | 1,078 (87.2%)                            | 251 (86.6%)                                    | 0.762           |
|                                   | No                              | 158 (12.8%)                              | 39 (13.4%)                                     |                 |
| Parity                            | Primiparous                     | 739 (59.8%)                              | 180 (62.1%)                                    | 0.476           |
|                                   | Multiparous                     | 497 (40.2%)                              | 110 (37.9%)                                    |                 |
| History of SA                     | Yes                             | 90 (7.3%)                                | 32 (11.0%)                                     | 0.034*          |
|                                   | No                              | 1,146 (92.7%)                            | 258 (89.0%)                                    |                 |
| Preconception care                | Yes                             | 786 (63.6%)                              | 144 (49.7%)                                    | <0.001*         |
|                                   | Not                             | 450 (36.4%)                              | 146 (50.3%)                                    |                 |

BMI, body mass index; SA, spontaneous abortion.

\*Bold values represent statistically significant results ( $p < 0.05$ ).

potentially modifiable factors can form the basis for clinical or preventive interventions in pregnancy management. However, given that most miscarriages occur in the early weeks of pregnancy, sometimes even before a woman knows she is pregnant, addressing these factors at this stage may be too late for any intervention or treatment to influence the outcome. This highlights the need to identify high-risk individuals before

conception to implement appropriate preventive measures aimed at reducing the risk (32).

We assessed the socio-demographic and clinical risk factors for first-trimester spontaneous abortion. Our results showed that the odds of first-trimester SA were higher in women over 35 years of age compared to women under 35 [[OR] = 2.02, 95% [CI] = 1.49–2.75]. This finding is consistent with previous


TABLE 3 Univariable and multivariable logistic regression for all candidate variables.

| Factors                           | Favorable outcome (%)<br>( <i>n</i> = 1,236) | SA in the first trimester (%)<br>( <i>n</i> = 290) | Univariable analysis [OR (95% CI)] | <i>p</i> -value | Multivariable analysis [OR (95% CI)] | <i>p</i> -value |
|-----------------------------------|----------------------------------------------|----------------------------------------------------|------------------------------------|-----------------|--------------------------------------|-----------------|
| Age of the woman (years): >35     | 15.9 (197/1,236)                             | 26.9 (78/290)                                      | 1.94 (1.44–2.62)                   | 0.001           | 2.02 (1.49–2.75)                     | 0.001           |
| Ethnicity                         |                                              |                                                    |                                    |                 |                                      |                 |
| Kazakh                            | 48.0 (593/1,236)                             | 47.9 (139/290)                                     | 1.14 (0.75–1.72)                   | 0.541           | –                                    | –               |
| Russian                           | 38.7 (478/1,236)                             | 40.4 (117/290)                                     | 1.19 (0.78–1.81)                   | 0.423           |                                      |                 |
| Other                             | 13.3 (165/1,236)                             | 11.7 (34/290)                                      | 1.00                               | –               |                                      |                 |
| Residency:                        |                                              |                                                    |                                    |                 |                                      |                 |
| Urban                             | 76.1 (941/1,236)                             | 79.7 (231/290)                                     | 1.23 (0.89–1.68)                   | 0.201           | –                                    | –               |
| Rural                             | 23.9 (295/1,236)                             | 20.3 (59/290)                                      | 1.00                               | –               |                                      |                 |
| Social status                     |                                              |                                                    |                                    |                 |                                      |                 |
| Employed, civil servant           | 5.7 (70/1,236)                               | 5.9 (17/290)                                       | 0.99 (0.57–1.75)                   | 0.986           | –                                    | –               |
| Employed, public sector worker    | 7.4 (92/1,236)                               | 6.6 (19/290)                                       | 0.85 (0.50–1.44)                   | 0.536           |                                      |                 |
| Employed, private sector worker   | 3.9 (48/1,236)                               | 3.8 (11/290)                                       | 0.94 (0.47–1.86)                   | 0.856           |                                      |                 |
| Employed, self-employed           | 1.7 (21/1,236)                               | 2.1 (6/290)                                        | 1.17 (0.46–2.96)                   | 0.739           |                                      |                 |
| Employed, laborer                 | 0.6 (8/1,236)                                | 0.3 (1/290)                                        | 0.51 (0.06–4.13)                   | 0.530           |                                      |                 |
| Employed, other                   | 7.4 (91/1,236)                               | 7.2 (21/290)                                       | 0.95 (0.57–1.58)                   | 0.830           |                                      |                 |
| Convicted                         | 0.3 (4/1,236)                                | 0 (0/290)                                          | 0.45 (0.02–8.48)                   | 0.597           |                                      |                 |
| Other                             | 44.4 (549/1,236)                             | 46.2 (134/290)                                     | 1.00                               | –               | –                                    | –               |
| Unemployed                        | 1.5 (19/1,236)                               | 1.4 (4/290)                                        | 0.86 (0.29–2.58)                   | 0.791           |                                      |                 |
| Housewife                         | 27.0 (334/1,236)                             | 26.6 (77/290)                                      | 0.94 (0.69–1.29)                   | 0.720           |                                      |                 |
| Education                         | 5.7 (70/1,236)                               | 5.9 (17/290)                                       | 0.99 (0.57–1.75)                   | 0.986           |                                      |                 |
| Secondary education               | 14.3 (177/1,236)                             | 13.8 (40/290)                                      | 0.96 (0.66–1.39)                   | 0.817           |                                      |                 |
| Higher than secondary education   | 85.7 (1,059/1,236)                           | 86.2 (250/290)                                     | 1.00                               | 1.00            |                                      |                 |
| Marital status                    |                                              |                                                    |                                    |                 |                                      |                 |
| Married                           | 83.2 (1,028/1,236)                           | 82.4 (239/1,236)                                   | 0.95 (0.68–1.33)                   | 0.757           | –                                    | –               |
| Not married                       | 16.8 (208/1,236)                             | 17.6 (51/1,236)                                    | 1.00                               |                 |                                      |                 |
| BMI (kg/m <sup>2</sup> ): Obesity | 5.3 (66/1,236)                               | 9.3 (27/290)                                       | 1.82 (1.14–2.90)                   | 0.012           | 1.81 (1.12–2.91)                     | 0.015           |
| Type 1 or Type 2 Diabetes         | 2.8 (34/1,236)                               | 5.22 (15/290)                                      | 1.93 (1.04–3.59)                   | 0.038           | –                                    | –               |
| Arterial Hypertension             | 3.3 (41/1,236)                               | 5.22 (15/290)                                      | 1.59 (0.87–2.91)                   | 0.134           | –                                    | –               |
| Hypothyroidism                    | 1.7 (21/1,236)                               | 1.7 (5/290)                                        | 1.02 (0.38–2.72)                   | 0.976           | –                                    | –               |
| Deficiency anemia                 | 25.1 (310/1,236)                             | 24.8 (72/290)                                      | 0.99 (0.73–1.32)                   | 0.929           | –                                    | –               |
| Allergies                         | 4.1 (51/1,236)                               | 5.9 (17/290)                                       | 1.45 (0.82–2.54)                   | 0.200           | –                                    | –               |
| Pre-pregnancy health insurance    | 87.2 (1,078/1,236)                           | 86.6 (251/290)                                     | 0.94 (0.65–1.37)                   | 0.761           | –                                    | –               |
| Parity                            |                                              |                                                    |                                    |                 |                                      |                 |
| Primiparous                       | 59.8 (739/1,236)                             | 62.1 (180/290)                                     | 1.10 (0.85–1.43)                   | 0.475           | –                                    | –               |
| Multiparous                       | 40.2 (497/1,236)                             | 37.9 (110/290)                                     | 1.00                               | –               |                                      |                 |
| History of SA                     | 7.3 (90/1,236)                               | 11.0 (32/290)                                      | 1.58 (1.03–2.42)                   | 0.035           | 1.57 (1.01–2.43)                     | 0.043           |
| Preconception Care                | 63.6 (786/1,236)                             | 49.7 (144/290)                                     | 0.57 (0.44–0.73)                   | 0.001           | 0.58 (0.45–0.75)                     | 0.001           |

BMI, body mass index; SA, spontaneous abortion; OR, odds ratio; CI, confidence interval. The univariable analysis included all variables from Table 2. The multivariable analysis included only variables retained in the final adjusted model; “–” indicates that the variable was not included. The outcome is coded as SA = 1.

research (33–35). The decline in reproductive potential in older women may be attributed to reduced oocyte quality, changes in the endometrium, and altered progesterone levels (36). Although age is a non-modifiable risk factor at the individual level, especially in cases of unintended pregnancies, it can be considered a modifiable risk factor at the population level. Including this factor in risk assessments can help prevent underestimating the risk of miscarriage in this population.

Furthermore, a strong negative impact of high BMI on pregnancy outcomes was observed [[OR] = 1.81, 95% [CI] = 1.12–2.91]. The risk of SA among women with obesity in our study aligns with findings from previous studies (37, 38). Possible mechanisms include poor endometrial receptivity, a pro-inflammatory state caused by cytokines, hormonal imbalances, or impaired blood supply to the endometrium and placenta (39–41). Further research is needed to confirm these



theories. Early weight loss interventions may reduce the risk of SA in obese patients. Public health policies targeting obesity prevention can significantly improve women’s reproductive outcomes (42). Such interventions should adopt a multisectoral approach addressing social, structural, economic, and environmental factors. Raising awareness among women about the importance of lifestyle changes and modifiable risk factors is necessary for reducing the overall risk of SA and other adverse maternal and infant health outcomes (12, 43–45).

We also found that women with a history of SA had an increased risk of first-trimester SA [[OR] = 1.57, 95% [CI] = 1.01–2.43], consistent with prior research indicating a significantly higher risk in this group (33). While previous pregnancy loss is a non-modifiable factor, including it in risk



assessments can prevent underestimating the risk in these populations.

An important finding of our study was that women who received preconception care had a lower risk of first-trimester SA [[OR] = 0.58, 95% [CI] = 0.45–0.75] compared to those who did not receive preconception care. This reinforces the conclusion that preconception care improves pregnancy outcomes (46). The preconception period offers a critical window for intervention, empowering women at higher risk of SA to address modifiable risk factors before conception (47). Potential mechanisms for reducing the risk of early pregnancy loss include optimizing chronic disease management, correcting nutritional deficiencies, normalizing weight, reducing harmful exposures, preventing infections, and taking folic acid, which aligns with meta-analysis findings (18, 48). These all highlight the need for national initiatives to improve the health of women of childbearing age before planned pregnancies. In Kazakhstan, the importance of preconception care has been recognized with the implementation of a new clinical protocol in 2023, providing comprehensive guidelines for healthcare providers to ensure its effective delivery (49). This protocol emphasizes the screening for chronic diseases, nutritional deficiencies, lifestyle modifications, vaccination, psychosocial support, counseling, and education to prepare couples for the emotional and physical challenges of pregnancy and parenthood.

The primary strength of this study is that, to the best of our knowledge, it is the first to investigate preconception care as a factor influencing first-trimester SA in women in Kazakhstan. Similar studies have been conducted in high-income countries with advanced healthcare systems, such as the United States and the United Kingdom, focusing on risk factors for early pregnancy loss and the broader implications of miscarriage (50, 51). Our study, conducted in Kazakhstan, applies to other Central Asian countries due to similar healthcare systems and their classification as middle-income economies. The analysis included a large sample of 1,526 women, which strengthens the statistical power and reliability of the results. These findings guide clinical counseling on spontaneous abortion risks and the public health benefits of preconception care. The study shows that preconception care protects against early pregnancy loss, providing clear support for clinical practice and public health efforts to improve pregnancy outcomes. This makes the results relevant for shaping policies and allocating resources. Integrating preconception care into routine health services, especially in low- and middle-income countries with limited access to such interventions, addresses modifiable risk factors like obesity and advanced maternal age. Such efforts improve maternal and neonatal health, lower healthcare costs, and promote health equity.

In addition to the factors identified in our analysis, long-term changes in the reproductive system, including hormonal disturbances, may also influence pregnancy outcomes. Emerging data suggest that some of these changes may be associated with prior COVID-19 infection, potentially affecting menstrual cycle regularity, ovulatory function, and endometrial receptivity (52–54). Although the clinical significance of these observations

requires further clarification, including such variables in future studies will enhance understanding of the mechanisms underlying adverse pregnancy outcomes and support the development of more precise preventive strategies.

From a public health perspective, the study has several limitations. First, selection bias is possible: the data were obtained from a single regional hospital with two perinatal centers that admit medium- and high-risk patients, and records with incomplete data were excluded. Second, information bias is possible: data and outcomes were extracted from an electronic medical database, and several covariates were missing, such as income, smoking, alcohol use, gestational age, antenatal complications, use of assisted reproductive technologies, or male factors (55, 56). Third, residual bias remains, since assignment to preconception care was not randomized, and exclusion of records with missing data may introduce selection bias if missingness is related to both exposure and outcome. The study period coincided with COVID-19 waves, which could have affected access to care, health-seeking behavior, and the structure of recorded cases. The moderate discriminative ability of the model (ROC-AUC  $\approx$  0.62) suggests that the results are most useful for assessing associations and planning prevention at the population level rather than for individual risk prediction. In addition, binary dichotomization of the outcome does not capture the severity of perinatal outcomes among live births and excludes late fetal or neonatal deaths. Future research should apply composite or ordinal perinatal endpoints. Finally, the reliance on retrospective data from medical records may underestimate the true prevalence of risk factors, particularly for socially sensitive or less commonly documented variables such as lifestyle behaviors or preconception care practices.

## 5 Conclusion

In our study, we evaluated associations between candidate factors and spontaneous abortion in the first trimester. Age >35 years, obesity, and a history of spontaneous abortion were associated with higher odds of early pregnancy loss, whereas preconception care was associated with lower odds. These findings expand the understanding of preventive measures taken before pregnancy in relation to adverse outcomes in the first trimester. In clinical practice, they may support risk stratification and counseling of women who could benefit from targeted preventive or therapeutic interventions before conception. From a public health perspective, the observed association with preconception care should be interpreted as associative rather than causal, and further prospective studies are needed before drawing conclusions about its impact on maternal and neonatal outcomes.

## Data availability statement

The data analyzed in this study is subject to the following licenses/restrictions: data is unavailable due to ethical and legal

restrictions. Requests to access these datasets should be directed to the Republican state enterprise on the right of economic management "Republican Center for Healthcare Development" of the Ministry of Health of the Republic of Kazakhstan, office@nrchd.kz

## Ethics statement

The studies involving humans were approved by Institutional Review Board of Karaganda Medical University (protocol code 22, dated 07.02.2022). The studies were conducted in accordance with the local legislation and institutional requirements. Written informed consent for participation was not required from the participants or the participants' legal guardians/next of kin in accordance with the national legislation and institutional requirements. Informed consent was obtained from all subjects involved in the study.

## Author contributions

YP: Conceptualization, Investigation, Writing – original draft, Methodology, Formal analysis, Validation, Writing – review & editing, Data curation. LS: Formal analysis, Supervision, Conceptualization, Methodology, Data curation, Writing – original draft, Writing – review & editing, Investigation. DK: Writing – review & editing, Conceptualization. YK: Investigation, Software, Methodology, Writing – original draft, Formal analysis, Data curation, Validation. ZA: Writing – original draft, Data curation. RS: Writing – review & editing, Writing – original draft, Supervision. OZ: Writing – original draft, Writing – review & editing.

## Funding

The author(s) declare that financial support was received for the research and/or publication of this article. The article processing charge is covered by Karaganda Medical University (Karaganda, Kazakhstan).

## References

1. World Health Organization. International classification of diseases (ICD). ICD-10 Version. (2019). Available online at: <https://icd.who.int/browse10/2019/en> (Accessed January 17, 2024).
2. Alves C, Jenkins SM, Rapp A. Early pregnancy loss (spontaneous abortion). *StatPearls*. Tampa/St. Petersburg, FL: StatPearls Publishing (2023). Available online at: <https://www.ncbi.nlm.nih.gov/books/NBK560521/> (Accessed January 17, 2024).
3. Committee on Practice Bulletins—Gynecology. The American college of obstetricians and gynecologists practice bulletin no. 150. Early pregnancy loss. *Obstet Gynecol*. (2015) 125:1258–67. doi: 10.1097/01.AOG.0000465191.27155.25
4. Strumpf E, Lang A, Austin N, Derksen SA, Bolton JM, Brownell MD, et al. Prevalence and clinical, social, and health care predictors of miscarriage. *BMC Pregnancy Childbirth*. (2021) 21:1–9. doi: 10.1186/S12884-021-03682-Z/TABLES/2
5. Hardy K, Hardy PJ. 1st Trimester miscarriage: four decades of study. *Transl Pediatr*. (2015) 4:189. doi: 10.3978/J.ISSN.2224-4336.2015.03.05
6. Cohain JS, Buxbaum RE, Mankuta D. Spontaneous first trimester miscarriage rates per woman among parous women with 1 or more pregnancies of 24 weeks or more. *BMC Pregnancy Childbirth*. (2017) 17:1–7. doi: 10.1186/S12884-017-1620-1/TABLES/3
7. Blushinova AN, Shalgumbayeva GM, Rakhmetova VS, Ryspaeva ZA, Aldabekova GU. Abortion practices among women in Kazakhstan: cross-sectional study. *Nauka I Zdravookhranenie (Science & Healthcare)*. (2024) 26(2):76–81. doi: 10.34689/SH.2024.26.2.010
8. Sakko Y, Turesheva A, Gaipov A, Aimagambetova G, Ukybassova T, Marat A, et al. Epidemiology of spontaneous pregnancy loss in Kazakhstan: a national population-based cohort analysis during 2014–2019 using the national electronic healthcare system. *Acta Obstet Gynecol Scand*. (2023) 102:1682. doi: 10.1111/AOGS.14669
9. Pendina AA, Krapivin MI, Chiryaeva OG, Petrova LI, Pashkova EP, Golubeva A V, et al. Chromosomal abnormalities in miscarriages and maternal age: new insights from the study of 7118 cases. *Cells*. (2024) 14:8. doi: 10.3390/CELLS14010008

## Acknowledgments

We acknowledge the administration of Karaganda Medical University for supporting the study.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Generative AI statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

Any alternative text (alt text) provided alongside figures in this article has been generated by Frontiers with the support of artificial intelligence and reasonable efforts have been made to ensure accuracy, including review by the authors wherever possible. If you identify any issues, please contact us.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgwh.2025.1615983/full#supplementary-material>

10. Kim YJ, Lee JE, Kim SH, Shim SS, Cha DH. Maternal age-specific rates of fetal chromosomal abnormalities in Korean pregnant women of advanced maternal age. *Obstet Gynecol Sci.* (2013) 56:160. doi: 10.5468/OGS.2013.56.3.160
11. Zhanseitova Advisor A, Crape B, Astana MB. *Spontaneous Abortion in Astana: Associated Factors, 2015–2017*. Astana: Nazarbayev University School of Medicine (2018). Available online at: <http://nur.nu.edu.kz/handle/123456789/3314> (Accessed January 17, 2024).
12. Metwally M, Ong KJ, Ledger WL, Li TC. Does high body mass index increase the risk of miscarriage after spontaneous and assisted conception? A meta-analysis of the evidence. *Fertil Steril.* (2008) 90:714–26. doi: 10.1016/J.FERTNSTERT.2007.07.1290
13. Muhammad T, Wan Y, Lv Y, Li H, Naushad W, Chan WY, et al. Maternal obesity: a potential disruptor of female fertility and current interventions to reduce associated risks. *Obes Rev.* (2023) 24:e13603. doi: 10.1111/OBR.13603
14. Sybäck K, Hartmann B, Kirchengast S. Maternal prepregnancy obesity affects foetal growth, birth outcome, mode of delivery, and miscarriage rate in Austrian women. *Int J Environ Res Public Health.* (2023) 20:4139. doi: 10.3390/IJERPH20054139
15. Huan Z, Yongping L, Lu L, Min Z, Xingzhi C, Yulong Q, et al. Maternal pre-pregnancy risk factors for miscarriage from a prevention perspective: a cohort study in China. *Eur J Obstet Gynecol.* (2016) 206:57–63. doi: 10.1016/j.ejogrb.2016.07.514
16. Yang J, Wang Y, Wang XY, Zhao YY, Wang J, Zhao YY. Adverse pregnancy outcomes of patients with history of first-trimester recurrent spontaneous abortion. *Biomed Res Int.* (2017) 2017:4359424. doi: 10.1155/2017/4359424
17. Sun H, Mao J, Su X, Du Q. Impact of spontaneous abortion history and induced abortion history on perinatal outcomes of singleton pregnancies. *BMC Public Health.* (2023) 23:1–10. doi: 10.1186/S12889-023-17264-5/TABLES/5
18. Magnus MC, Hockey RL, Häberg SE, Mishra GD. Pre-pregnancy lifestyle characteristics and risk of miscarriage: the Australian longitudinal study on women's health. *BMC Pregnancy Childbirth.* (2022) 22:1–10. doi: 10.1186/S12884-022-04482-9/TABLES/3
19. Al-Alami Z, Abu-Huwajir R, Hamadneh S, Taybeh E. Understanding miscarriage prevalence and risk factors: insights from women in Jordan. *Medicina (B Aires).* (2024) 60:1044. doi: 10.3390/MEDICINA60071044
20. Feodor Nilsson S, Andersen PK, Strandberg-Larsen K, Nybo Andersen AM. Risk factors for miscarriage from a prevention perspective: a nationwide follow-up study. *BJOG.* (2014) 121:1375–85. doi: 10.1111/1471-0528.12694
21. *Preconception Care*. Geneva: World Health Organization (WHO) (2017). Available online at: <https://iris.who.int/handle/10665/205637> (Accessed January 18, 2024).
22. Xu J, Li X, Zhou Q. Nationwide-free preconception care strategy: experience from China. *Front Public Health.* (2022) 10:934983. doi: 10.3389/FPUBH.2022.934983/BIBTEX
23. Dean S V, Lassi ZS, Imam AM, Bhutta ZA. Preconception care: closing the gap in the continuum of care to accelerate improvements in maternal, newborn and child health. *Reprod Health.* (2014) 11:1–8. doi: 10.1186/1742-4755-11-S3-S1/FIGURES/3
24. Poix S, Elmusharaf K. Investigating the pathways from preconception care to preventing maternal, perinatal and child mortality: a scoping review and causal loop diagram. *Prev Med Rep.* (2023) 34:102274. doi: 10.1016/J.PMEDR.2023.102274
25. Clinical Protocol for the Diagnosis and Treatment of Pregnancy Loss Approved by the Unified Commission for the Quality of Medical Services of the Ministry of Health of the Republic of Kazakhstan on July 28, 2023, Protocol No. 185. Available online at: <https://nrchd.kz/ru/2017-03-12-10-51-13/klinicheskie-protokoly> (Accessed January 18, 2024).
26. *Early Pregnancy Loss*. Washington, DC: American College of Obstetricians and Gynecologists (ACOG) (2018). Available online at: <https://www.acog.org/clinical/clinical-guidance/practice-bulletin/articles/2018/11/early-pregnancy-loss> (Accessed January 18, 2024).
27. On Approval of the Standard for Organizing Obstetric and Gynecological Care in the Republic of Kazakhstan Order of the Minister of Health of the Republic of Kazakhstan No-92, August 26, 2021. Available online at: <https://adilet.zan.kz/kaz/docs/V2100024131> (Accessed January 18, 2024).
28. *Obesity: Preventing and Managing the Global Epidemic: Report of a WHO Consultation*. Geneva: World Health Organization (WHO) (2000). Available online at: <https://iris.who.int/handle/10665/42330> (Accessed January 18, 2024).
29. Choo V. WHO Reassesses appropriate body-mass index for Asian populations. *Lancet.* (2002) 360:235. doi: 10.1016/S0140-6736(02)09512-0
30. Clinical Protocol for the Diagnosis and Treatment of Arterial Hypertension Approved by the Unified Commission for the Quality of Medical Services of the Ministry of Health of the Republic of Kazakhstan on October 3, 2019, Protocol No. 74. Available online at: [https://online.zakon.kz/Document/?doc\\_id=37136070&pos=3;44#pos=3;44](https://online.zakon.kz/Document/?doc_id=37136070&pos=3;44#pos=3;44) (Accessed January 18, 2024).
31. World Health Organization. WHO Recommendations on antenatal care for a positive pregnancy experience. *WHO Recommendations on Antenatal Care for a Positive Pregnancy Experience*. Geneva: World Health Organization (WHO) (2016). Available online at: <https://www.ncbi.nlm.nih.gov/books/NBK409108/> (Accessed January 18, 2024).
32. Taizhanova D, Zubkov DV, Komlichenko EV, Magalov I, Sorokina MA, Bepalova NV, et al. The possibilities of adverse pregnancy outcomes predicting based on laboratory markers of reproductive losses. *Med Ecol.* (2025) 0:77–84. doi: 10.59598/ME-2305-6045-2024-113-4-77-84
33. Magnus MC, Wilcox AJ, Morken NH, Weinberg CR, Häberg SE. Role of maternal age and pregnancy history in risk of miscarriage: prospective register based study. *Br Med J.* (2019) 364:869. doi: 10.1136/BMJ.L869
34. Cavazos-Reh PA, Krauss MJ, Spitznagel EL, Bommarito K, Madden T, Olsen MA, et al. Maternal age and risk of labor and delivery complications. *Matern Child Health J.* (2015) 19:1202–11. doi: 10.1007/S10995-014-1624-7
35. Koo YJ, Ryu HM, Yang JH, Lim JH, Lee JE, Kim MY, et al. Pregnancy outcomes according to increasing maternal age. *Taiwan J Obstet Gynecol.* (2012) 51:60–5. doi: 10.1016/J.TJOG.2012.01.012
36. Soler A, Morales C, Mademont-Soler I, Margarit E, Borrell A, Borobio V, et al. Overview of chromosome abnormalities in first trimester miscarriages: a series of 1,011 consecutive chorionic villi sample karyotypes. *Cytogenet Genome Res.* (2017) 152:81–9. doi: 10.1159/000477707
37. Ghimire PR, Akombi-Inyang BJ, Tannous C, Agho KE. Association between obesity and miscarriage among women of reproductive age in Nepal. *PLoS One.* (2020) 15:e0236435. doi: 10.1371/JOURNAL.PONE.0236435
38. Silvestris E, de Pergola G, Rosania R, Loverro G. Obesity as disruptor of the female fertility. *Reprod Biol Endocrinol.* (2013) 16:1–13. doi: 10.1186/S12958-018-0336-Z
39. Snider AP, Wood JR. Obesity induces ovarian inflammation and reduces oocyte quality. *Reproduction.* (2019) 158:R79–90. doi: 10.1530/REP-18-0583
40. Metwally M, Preece R, Thomas J, Ledger W, Li TC. A proteomic analysis of the endometrium in obese and overweight women with recurrent miscarriage: preliminary evidence for an endometrial defect. *Reprod Biol Endocrinol.* (2014) 12:75. doi: 10.1186/1477-7827-12-75
41. Macroscopic and Microscopic Morphological Features of The Placenta Associated with Preeclampsia and Chronic Hypoxic Damage of the Fetus | Muhammad | Medicine and ecology. (Accessed January 22, 2024).
42. Kalantari E, Tajvar M, Naderimaghani S, Takian A. Maternal obesity management: a narrative literature review of health policies. *BMC Womens Health.* (2024) 24:1–12. doi: 10.1186/S12905-024-03342-2/TABLES/4
43. Belan M, Harnois-Leblanc S, Laferrère B, Baillargeon JP. Optimizing reproductive health in women with obesity and infertility. *CMAJ.* (2018) 190:E742. doi: 10.1503/CMAJ.171233
44. Nukeshtayeva K, Kayupova G, Yerdessov N, Bolatova Z, Zhamantayev O, Turmukhambetova A. Factors associated with maternal mortality in Kazakhstan: a pre- and during-pandemic comparison. *Front Public Health.* (2024) 12:1337564. doi: 10.3389/FPUBH.2024.1337564
45. Yerdessov N, Zhamantayev O, Bolatova Z, Nukeshtayeva K, Kayupova G, Turmukhambetova A. Infant mortality trends and determinants in Kazakhstan. *Children.* (2023) 10:923. doi: 10.3390/CHILDREN10060923
46. Timraz M, Abu-Hamad B, Ibaid A. Does preconception care make a difference to pregnancy outcomes? A quasi-experimental, mixed-methods study. *Lancet.* (2022) 399:S44. doi: 10.1016/s0140-6736(22)01179-5
47. Benedetto C, Borella F, Divakar H, O'Riordan SL, Mazzoli M, Hanson M, et al. FIGO preconception checklist: preconception care for mother and baby. *Int J Gynaecol Obstet.* (2024) 165:1–8. doi: 10.1002/IJGO.15446
48. Wahabi HA, Fayed A, Esmail S, Elmorshedy H, Titi MA, Amer YS, et al. Systematic review and meta-analysis of the effectiveness of pre-pregnancy care for women with diabetes for improving maternal and perinatal outcomes. *PLoS One.* (2020) 15(8):e0237571. doi: 10.1371/journal.pone.0237571
49. Clinical Protocol for Diagnosis and Treatment: Preconception Care, approved by the Joint Kat Commission on the Quality of Medical Services of the Ministry of Health of the Republic of Kazakhstan, dated July 28, 2023, Protocol No. 185. Available online at: <https://nrchd.kz/ru/2017-03-12-10-51-13/klinicheskie-protokoly> (Accessed January 20, 2024).
50. Gracia CR, Sammel MD, Chittams J, Hummel AC, Shaunik A, Barnhart KT. Risk factors for spontaneous abortion in early symptomatic first-trimester pregnancies. *Obstet Gynecol.* (2005) 106:993–9. doi: 10.1097/01.AOG.0000183604.09922.E0
51. Quenby S, Gallos ID, Dhillon-Smith RK, Podsek M, Stephenson MD, Fisher J, et al. Miscarriage matters: the epidemiological, physical, psychological, and economic costs of early pregnancy loss. *Lancet.* (2021) 397:1658–67. doi: 10.1016/S0140-6736(21)00682-6/ATTACHMENT/CD1B82A0-EB07-4548-B733-4399095F3D6E/MMC1.PDF
52. Lebar V, Laganà AS, Chiantera V, Kunič T, Lukanović D. The effect of COVID-19 on the menstrual cycle: a systematic review. *J Clin Med.* (2022) 11:3800. doi: 10.3390/jcm11133800

53. Ghaemi M, Hantoushzadeh S, Shafiee A, Gargari OK, Fathi H, Eshraghi N, et al. The effect of COVID-19 and COVID-19 vaccination on serum anti-mullerian hormone: a systematic review and meta-analysis. *Immun Inflamm Dis.* (2024) 12: e1136. doi: 10.1002/iid3.1136
54. Huang J, Liu Y, Wang J, Xu D, Huang Z, Li M, et al. Pregnancy outcomes after frozen-thawed embryo transfer in women with COVID-19 history: a prospective cohort study. *J Med Virol.* (2024) 96:e29377. doi: 10.1002/jmv.29377
55. Kleinhaus K, Perrin M, Friedlander Y, Paltiel O, Malaspina D, Harlap S. Paternal age and spontaneous abortion. *Obstet Gynecol.* (2006) 108:369–77. doi: 10.1097/01.AOG.0000224606.26514.3A
56. Robinson L, Gallos ID, Conner SJ, Rajkhowa M, Miller D, Lewis S, et al. The effect of sperm DNA fragmentation on miscarriage rates: a systematic review and meta-analysis. *Hum Reprod.* (2012) 27:2908–17. doi: 10.1093/HUMREP/DES261