



OPEN ACCESS

APPROVED BY
Frontiers Editorial Office,
Frontiers Media SA, Switzerland

*CORRESPONDENCE

Tuerganaili Aji
✉ tuergan78@asina.com

RECEIVED 10 October 2025

ACCEPTED 10 October 2025

PUBLISHED 24 October 2025

CITATION

Abulizi A, Tuergan T, Shalayiadang P, Zhang C, Zhang R, Jiang T, Guo Q, Wang H, Li L, Lin R, Shao Y and Aji T (2025) Correction: Hepatic alveolar echinococcosis infection induces a decrease in NK cell function through high expression of NKG2A in patients. *Front. Immunol.* 16:1638960. doi: 10.3389/fimmu.2025.1638960

COPYRIGHT

© 2025 Abulizi, Tuergan, Shalayiadang, Zhang, Zhang, Jiang, Guo, Wang, Li, Lin, Shao and Aji. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Correction: Hepatic alveolar echinococcosis infection induces a decrease in NK cell function through high expression of NKG2A in patients

Abuduaini Abulizi^{1,2}, Talaiti Tuergan^{1,2}, Paizula Shalayiadang^{1,2}, Chuanshan Zhang^{2,3}, Ruiqing Zhang^{1,2}, Tiemin Jiang^{1,2}, Qiang Guo^{1,2}, Hui Wang^{2,3}, Liang Li^{2,3}, Renyong Lin^{2,3}, Yingmei Shao^{1,2} and Tuerganaili Aji^{1,2*}

¹Hepatobiliary & Hydatid Disease Department, Digestive & Vascular Surgery Center, First Affiliated Hospital of Xinjiang Medical University, State Key Laboratory of Pathogenesis, Prevention and Treatment of High Incidence Diseases in Central Asia, Urumqi, China, ²World Health Organization (WHO) Collaborating Center on Prevention and Management of Echinococcosis, Clinical Medicine Institute, First Affiliated Hospital of Xinjiang Medical University, Urumqi, China, ³State Key Laboratory on Pathogenesis Prevention & Treatment of High Incidence Diseases in Central Asia, Xinjiang Medical University, Urumqi, China

KEYWORDS

hepatic alveolar echinococcosis, NK cell, NKG2A, immune exhaustion, IFN- γ

A Correction on

Hepatic alveolar echinococcosis infection induces a decrease in NK cell function through high expression of NKG2A in patients

By Abulizi A, Tuergan T, Shalayiadang P, Zhang C, Zhang R, Jiang T, Guo Q, Wang H, Li L, Lin R, Shao Y and Aji T (2025). *Front. Immunol.* 16:1563248. doi: 10.3389/fimmu.2025.1563248

Author “Talaiti Tuergan” was erroneously assigned as equal contributing author. The original version of this article has been updated.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.