



OPEN ACCESS

EDITED AND REVIEWED BY
Pavel Lindberg,
INSERM U1266 Institut de Psychiatrie et
Neurosciences de Paris, France

*CORRESPONDENCE
Adriana Bastos Conforto
✉ adriana.conforto@einstein.br

RECEIVED 26 September 2025
ACCEPTED 03 October 2025
PUBLISHED 17 October 2025

CITATION
Kroth JB, Handfas B, Rodrigues G, Zepeda F,
Oliveira MA, Wang DJJ, Azevedo Neto RMD,
Silva GS, Amaro E Jr, Sorinola IO and
Conforto AB (2025) Correction: Effects of
repetitive peripheral sensory stimulation in the
subacute and chronic phases after stroke:
study protocol for a pilot randomized trial.
Front. Neurol. 16:1713834.
doi: 10.3389/fneur.2025.1713834

COPYRIGHT
© 2025 Kroth, Handfas, Rodrigues, Zepeda,
Oliveira, Wang, Azevedo Neto, Silva, Amaro,
Sorinola and Conforto. This is an open-access
article distributed under the terms of the
[Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in
other forums is permitted, provided the
original author(s) and the copyright owner(s)
are credited and that the original publication
in this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Correction: Effects of repetitive peripheral sensory stimulation in the subacute and chronic phases after stroke: study protocol for a pilot randomized trial

Jéssica Borges Kroth¹, Benjamim Handfas¹, Glaucia Rodrigues¹,
Francisco Zepeda², Marco Aurélio Oliveira¹, Danny J. J. Wang³,
Raymundo Machado de Azevedo Neto¹, Gisele Sampaio Silva¹,
Edson Amaro Jr.¹, Isaac Olubunmi Sorinola⁴ and
Adriana Bastos Conforto^{1*}

¹Hospital Israelita Albert Einstein, São Paulo, Brazil, ²Biological Engineering Department, Massachusetts Institute of Technology, Boston, MA, United States, ³Department of Neurology, Keck School of Medicine, University of Southern California, Los Angeles, CA, United States, ⁴Department of Physiotherapy, King's College London, London, United Kingdom

KEYWORDS

sensory stimulation, stroke, rehabilitation, upper limb, nerve stimulation

A Correction on

Effects of repetitive peripheral sensory stimulation in the subacute and chronic phases after stroke: study protocol for a pilot randomized trial

by Kroth, J. B., Handfas, B., Rodrigues, G., Zepeda, F., Oliveira, M. A., Wang, D. J. J., Azevedo Neto, R. M. d., Silva, G. S., Amaro, E. Jr., Sorinola, I. O., and Conforto, A. B. (2022). *Front. Neurol.* 13:779128. doi: 10.3389/fneur.2022.779128

In the published article, there was an error in [Figure 1](#) as published. Three performance measures—Jebson-Taylor test, pinch strength, and grasp strength—were obtained at three time points: before repetitive peripheral sensory stimulation, after stimulation, and after training. The original figure showed only two time points. The corrected [Figure 1](#) and caption are provided below.

The original version of this article has been updated.

Generative AI statement

Any alternative text (alt text) provided alongside figures in this article has been generated by Frontiers with the support of artificial intelligence and reasonable efforts have been made to ensure accuracy, including review by the authors wherever possible. If you identify any issues, please contact us.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated

organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.




FIGURE 1
Experimental paradigm. NIH, national institutes of health; PHQ-9, patient health questionnaire-9; MMSE, mini-mental state exam; MRI, magnetic resonance imaging; RPSS, repetitive peripheral nerve sensory stimulation; JTT, Jebsen–Taylor test.