



OPEN ACCESS

EDITED AND REVIEWED BY

Maria A. Tikhonova,
Scientific Research Institute of Neurosciences
and Medicine, Russia

*CORRESPONDENCE

Xueqian Wang
✉ shirlyding@163.com
Qingguo Wang
✉ wangqg8558@163.com
Fafeng Cheng
✉ fafengcheng@gmail.com

[†]These authors have contributed equally to
this work and share first authorship

RECEIVED 23 November 2025

ACCEPTED 25 November 2025

PUBLISHED 04 December 2025

CITATION

Xu W, Ren B, Zhang Z, Chen C, Xu T, Liu S,
Ma C, Wang X, Wang Q and Cheng F (2025)
Correction: Network pharmacology analysis
reveals neuroprotective effects of the
Qin-Zhi-Zhu-Dan Formula in Alzheimer's
disease. *Front. Neurosci.* 19:1752611.
doi: 10.3389/fnins.2025.1752611

COPYRIGHT

© 2025 Xu, Ren, Zhang, Chen, Xu, Liu, Ma,
Wang, Wang and Cheng. This is an
open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](#). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that the
original publication in this journal is cited, in
accordance with accepted academic practice.
No use, distribution or reproduction is
permitted which does not comply with these
terms.

Correction: Network pharmacology analysis reveals neuroprotective effects of the Qin-Zhi-Zhu-Dan Formula in Alzheimer's disease

Wenxiu Xu^{1†}, Beida Ren^{2,3†}, Zehan Zhang¹, Congai Chen²,
Tian Xu¹, Shuling Liu¹, Chongyang Ma⁴, Xueqian Wang^{1*},
Qingguo Wang^{1*} and Fafeng Cheng^{1*}

¹School of Traditional Chinese Medicine, Beijing University of Chinese Medicine, Beijing, China,

²Dongzhimen Hospital, Beijing University of Chinese Medicine, Beijing, China, ³Institute for Brain
Disorders, Beijing University of Chinese Medicine, Beijing, China, ⁴School of Traditional Chinese
Medicine, Capital Medical University, Beijing, China

KEYWORDS

Alzheimer's disease, Qin-Zhi-Zhu-Dan Formula, TNF signaling pathway, network
pharmacology, inflammatory response

A Correction on

Network pharmacology analysis reveals neuroprotective effects of the
Qin-Zhi-Zhu-Dan Formula in Alzheimer's disease

by Xu, W., Ren, B., Zhang, Z., Chen, C., Xu, T., Liu, S., Ma, C., Wang, X., Wang, Q., and Cheng, F.
(2022). *Front. Neurosci.* 16:943400. doi: 10.3389/fnins.2022.943400

There was a mistake in [Figure 10B](#) as published. The higher magnification images in the
QZZD and DNZ columns were incorrectly swapped: the image labeled QZZD belonged to
the DNZ group, and the image labeled DNZ belonged to the QZZD group. The corrected
[Figure 10](#) appears below.

There was a mistake in [Figure 11C](#) as published. An image designated for the QZZD
group was incorrectly reused in both QZZD and DNZ panels. The corrected [Figure 11](#)
appears below.

The original version of this article has been updated.

Generative AI Statement

Any alternative text (alt text) provided alongside figures in this article has been
generated by Frontiers with the support of artificial intelligence and reasonable efforts have
been made to ensure accuracy, including review by the authors wherever possible. If you
identify any issues, please contact us.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily
represent those of their affiliated organizations, or those of the publisher, the editors and
the reviewers. Any product that may be evaluated in this article, or claim that may be made
by its manufacturer, is not guaranteed or endorsed by the publisher.




FIGURE 10
Effects of QZZD on neuronal death. **(A)** CA1 region of the hippocampus after HE staining [Scale bar: 75 and 25 μ m (insert), $n = 6$]. **(B)** CA1 region of the hippocampus after NeuN staining [Scale bar: 75 and 25 μ m (insert)]. **(C)** Number of NeuN-positive cells in CA1 region of the hippocampus. Values were expressed as means \pm SD, $n = 6$ in each group. ** $p < 0.01$, compared with the control group; ## $p < 0.01$, compared with the model group.




FIGURE 11

Effects of QZZD on expressions of Aβ, IL-6, NF-κBp65, TNFR1, p-ERK1/2 and ERK1/2 in hippocampal tissues of APP/PS1 double transgenic mice with dementia. **(A)** Western blotting analysis showing the protein expression levels of Aβ, IL-6, NF-κBp65, TNFR1, p-ERK1/2 and ERK1/2 in the hippocampus. **(B)** Quantitative analysis of Aβ, IL-6, NF-κBp65, TNFR1, p-ERK1/2, and ERK1/2 protein bands. Values are expressed as the mean ± SD, $n = 6$ in each group. ** $p < 0.01$, compared with the control group; # $p < 0.05$, ## $p < 0.01$, compared with the model group. **(C)** Immunohistochemistry analysis showing the protein expression levels of Aβ, TNFR1, NF-κBp65, and IL-6 in the hippocampus. **(D)** The average optical density of Aβ, TNFR1, NF-κBp65 and IL-6 in the hippocampus. Values are expressed as the mean ± SD, $n = 6$ for each group. ** $p < 0.01$, compared with the control group; # $p < 0.05$, ## $p < 0.01$, compared with the model group.