



OPEN ACCESS

EDITED BY

Lara Costantini,
University of Tuscia, Italy

REVIEWED BY

Rajesh Kanna Gopal,
Saveetha Dental College and Hospital,
SIMATS, India
Yoredy Sarmiento-Andrade,
Universidad Técnica Particular de Loja,
Ecuador

*CORRESPONDENCE

Adriyan Pramono
✉ adriyanpramono@fk.undip.ac.id

[†]These authors have contributed equally to this work and share first authorship

RECEIVED 17 June 2025

ACCEPTED 13 August 2025

PUBLISHED 08 September 2025

CITATION

Fadhillah FS, Arafat FB, Maharani N, Fadhillah J, Zariyah H, Sobirin MA, Tsutsui H, Wibowo A, Juniarto AZ and Pramono A (2025) Gut microbiota profile in adults with abdominal obesity and non-abdominal obesity: links to body composition and macronutrient intake in Semarang, Indonesia—a brief research report. *Front. Nutr.* 12:1648575. doi: 10.3389/fnut.2025.1648575

COPYRIGHT

© 2025 Fadhillah, Arafat, Maharani, Fadhillah, Zariyah, Sobirin, Tsutsui, Wibowo, Juniarto and Pramono. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Gut microbiota profile in adults with abdominal obesity and non-abdominal obesity: links to body composition and macronutrient intake in Semarang, Indonesia—a brief research report

Farhan Syafiq Fadhillah^{1†}, Fariz Budi Arafat^{2†}, Nani Maharani³, Jihan Fadhillah¹, Hainun Zariyah¹, Mochamad Ali Sobirin⁴, Hiroyuki Tsutsui⁵, Adi Wibowo⁶, Achmad Zulfa Juniarto⁷ and Adriyan Pramono^{1,8*}

¹Department of Nutrition Science, Faculty of Medicine, Diponegoro University, Semarang, Indonesia,

²Department of Statistics, Faculty of Sciences and Mathematics, Diponegoro University, Semarang,

Indonesia, ³Department of Pharmacology and Therapy, Faculty of Medicine, Diponegoro University,

Semarang, Indonesia, ⁴Department of Cardiology and Vascular Medicine, Faculty of Medicine, Diponegoro

University, Semarang, Indonesia, ⁵Department of Cardiovascular Medicine, Faculty of Medical Sciences,

Kyushu University, Fukuoka, Japan, ⁶Department of Informatics, Faculty of Sciences and Mathematics,

Diponegoro University, Semarang, Indonesia, ⁷Center of Biomedical Research, Diponegoro University,

Semarang, Indonesia, ⁸Center of Nutrition Research, Diponegoro University, Semarang, Indonesia

Background: Abdominal obesity is a significant risk factor for metabolic syndrome and cardiovascular diseases, with increasing evidence highlighting the role of gut microbiota in its development. In Indonesia, where 23.4% of adults are obese, few studies have examined the gut microbiota in relation to abdominal obesity, particularly in the context of unique dietary patterns. This pilot study investigated the gut microbiota composition in adults with abdominal obesity in Semarang, Indonesia, and its associations with body composition and macronutrient intake.

Methods: This cross-sectional study was conducted in Semarang and included 46 adults aged 20–50 years, categorized by abdominal obesity status (22 with abdominal obesity and 24 without). Anthropometric measurements, body composition, and dietary intake were assessed. Gut microbiota profiles were analyzed using 16S rRNA gene sequencing of fecal samples.

Findings: In the Semarang population, individuals with abdominal obesity had higher visceral fat ($12.32 \pm 3.44\%$ vs. $6.96 \pm 2.91\%$) compared to those without abdominal obesity. *Prevotella_9 copri* was positively associated with visceral fat ($r = 0.206$, $p = 0.169$), a finding that differs from studies conducted outside Indonesia, potentially showing the uniqueness of the profile.


Conclusion: The correlation of *Prevotella_9 copri* in subjects from Semarang, Indonesia, differs from findings in other studies, providing a potential unique gut microbiota profile in the Indonesian population and providing a platform for future studies to clarify these hypotheses. Larger longitudinal studies are needed to validate these findings and establish causality.

KEYWORDS

gut microbiota, abdominal obesity, microbial diversity, adults, dietary intake



- In Indonesia, where 23.4% of adults are obese, there is a lack of studies examining gut microbiota in the context of abdominal obesity, particularly given the unique dietary patterns.
- This study investigated the gut microbiota composition in adults with abdominal obesity in Semarang, Indonesia, and its associations with body composition and macronutrient intake



Cross-sectional study in Semarang, Indonesia


22 Abdominal obesity

24 Non-Abdominal Obesity



- 20-50 years old
- Without chronic diseases
- Non pregnant and breastfeeding

Measurements of
Anthropometric, Body Composition, Gut Microbiota from fecal



Heatmap analysis shown correlation with indicator of metabolic health

- *Prevotella_9 Copri* is positively associated with Visceral Fat, different from other study that are negatively associated
- These findings providing potential uniqueness of the profile of gut microbiota in Indonesia compared to other countries.
- Larger longitudinal studies are needed to validate these findings and establish causality.

GRAPHICAL ABSTRACT

1 Introduction

Obesity affected 16% of the global adult population in 2024, with developing countries such as Indonesia currently experiencing rising rates; 23.4% of Indonesian adults were obese in 2023, up from 21.8% in 2018 (1, 2). Jakarta, the capital region, has a prevalence of abdominal obesity of 31.8%. There are some interesting findings where the coastal districts have a lower prevalence (for example, in the Nusa Tenggara Timur Province, where majority of the population lives in the rural areas, the prevalence is 13.3%) (2–5). Indonesia has distinct geographical characteristics, from coastal to non-coastal, with one study showing an improvement in the health quality in the coastal area. A study from Geiger SJ et al. shows better reported self-health in population near the coastal area (6). However, the study has yet to clarify whether living in coastal area is associated with lower body mass index (BMI) and/or reduced obesity prevalence. While rapid urbanization, nutritional transition, and lifestyle changes may have contributed to these trends, other factors may also explain the difference between abdominal obesity and other factors, such as gut microbiota where different dietary intake can alter the gut microbiota composition, producing different nutritional metabolism (7), possibly explaining the uniqueness of Indonesian population, where predominantly coastal living is associated with lower BMI and overall lower abdominal obesity (8). With that perspective, the study that overviews the profile of gut microbiota in adults with abdominal obesity and comparing with non-abdominal obesity in

Indonesia is necessary, given the diversity of dietary intake in urban and rural areas of Indonesia (9).

BMI may not fully capture health risks from fat distribution; abdominal circumference better indicates visceral fat accumulation. Abdominal obesity's strong link to visceral fat and metabolic complications has prompted microbiota investigations. Visceral fat, more than subcutaneous fat, correlates with metabolic syndrome, insulin resistance, and cardiovascular disease (4). Recognizing these associations is critical for public health and clinical practice, as understanding fat distribution patterns can improve early interventions and reduce obesity complications. Additionally, research indicates that gut microbiota significantly influences metabolic health. Individuals with altered microbial communities show metabolic changes with increased calorie absorption and adiposity, suggesting the microbiome's potential as a metabolic biomarker and therapeutic target (10, 11). Differences in locations, such as coastal and non-coastal areas, could alter human gut microbiota through variations in dietary intake. Fiçıcılar's study in Samsun, Turkey, showed higher fish intake in coastal areas, while Bratlie et al.'s research in Norway found that fish consumption was correlated with higher Firmicutes and lower Bacteroidetes compared to controls (12, 13).

To bridge these insights, this pilot study integrates abdominal circumference measures with gut microbiota profiles, body composition, and macronutrient intake in Indonesian adults with abdominal obesity and non-abdominal obesity, conducted by a cross-sectional study in

Semarang, Indonesia, exploring the potential idea to benefit the population, researchers, and policymakers in precision nutrition.

2 Method

2.1 Study design

This cross-sectional study was conducted in northern Semarang, Indonesia, from June to December 2024, to examine the profile of the gut microbiota in adults with abdominal obesity and non-abdominal obesity. Semarang was selected for its rapid urbanization and evolving dietary patterns that reflect transitions in emerging economies (14). Data were collected from local community centers and health facilities across several sites. To minimize potential sources of bias, the data collection was conducted on the same day as fecal sampling, reducing the chance of fabricated results such as having dietary intake before or after the fecal collection. The study protocol was reviewed and approved by the Medical Ethics Committee of the Faculty of Medicine, Diponegoro University (322/EC/KEPK/FK-UNDIP/VI/2024).

2.2 Participants

The study enrolled adults aged 20–50 years residing in the northern part of Semarang, Indonesia, who were categorized based on their abdominal obesity status, with abdominal obesity defined as waist circumference of ≥ 90 cm (men) or ≥ 80 cm (women), according to the Indonesian cutoff (15). Exclusion criteria were chronic metabolic diseases, pregnancy, or breastfeeding. Participants were screened through questionnaires and clinical assessments. Recruitment occurred via community outreach, advertisements, and primary care providers. All participants provided informed consent per Declaration of Helsinki (16). This study enrolled a total of 46 participants: with 22 in abdominal obesity and 24 in non-abdominal obesity group. The sample size was limited by low participation during the one-month recruitment period in June and by resource constraints. The limited recruitment duration may introduce participation bias; however, as most participants were recruited from two clinical centers and given the nature of this pilot study, participation bias was not considered.

2.3 Study variables and measurements

2.3.1 Anthropometric and vital sign measurements

Participants body weight and height were measured using calibrated digital scales and stadiometers. These values were used to calculate BMI as kg/m^2 . In addition, waist and hip circumferences were measured at standardized anatomical landmarks using non-stretchable tape. Waist circumference was emphasized as a critical measure because it reflects visceral fat accumulation, which is strongly associated with metabolic risk factors (17). Additionally, we measured the systole and diastole of the participants blood pressure.

2.3.2 Body composition analysis

Body composition parameters, including fat mass, muscle mass, and visceral fat index, were evaluated using bioelectrical impedance analysis (BIA) (Tanita, Tokyo, Japan). BIA was selected because it is noninvasive, rapid, and cost-effective, with proven reliability compared to techniques such as dual-energy X-ray absorptiometry (DXA) (18).

2.3.3 Dietary intake

Macronutrient and energy intakes were assessed using a validated semi-quantitative food frequency questionnaire (SQ-FFQ) specifically designed for the Indonesian population. The SQ-FFQ was pre-tested for reliability and validity in similar cohorts to capture the frequency and portion sizes of commonly consumed foods (19), and it is chosen instead of other tools such as diet quality questionnaire (DQQ) because of the non-specific and generalized nature of DQQ itself (20).

2.3.4 Fecal DNA isolation and sequencing

Fecal samples were collected in sterile containers at -20°C during transport and processed at the laboratory. DNA was extracted from 200 mg fecal matter using QIAamp PowerFecal Pro DNA Kit (Qiagen, Hilden, Germany) per manufacturer's protocol. DNA quality and yield were verified by spectrophotometry and gel electrophoresis. The V4 region of bacterial 16S rRNA gene was sequenced using Illumina MiSeq platform (Illumina, San Diego, CA, USA). Studies have validated this method for characterizing gut microbial communities (21).


2.4 Statistical methods

Data analysis was performed using R programming language (version 4.5.0; R Core Team, Vienna, Austria), with microbiome data processed using the phyloseq package (version 1.51.0). Baseline data were summarized using descriptive statistics. Groups defined by abdominal circumference were compared using the Mann–Whitney U test for alpha diversity, while beta diversity was assessed using Bray–Curtis dissimilarity calculated with the vegan package (v2.6–10) in R. Non-metric multidimensional scaling (NMDS) and principal coordinates analysis (PCoA) were used to visualize beta diversity patterns. Group-level differences were statistically evaluated using PERMANOVA. The taxa abundance was analyzed using QIIME2 (QIIME Development Team, Tucson, AZ, USA) and DESeq2 (22). Spearman rank correlation was used to assess associations between gut microbiota profiles and clinical parameters. To account for multiple comparisons, *p*-values were adjusted using the Benjamini–Hochberg procedure (false discovery rate, FDR) (23). The Spearman correlation analysis showed species with $\geq 80\%$ abundance best visualized gut microbiota associations with biomarkers. A heatmap illustrated these correlations, enabling the interpretation of microbial patterns and inter-sample variability, as shown in other studies (24).

3 Results

3.1 Participants

A total of 46 participants were recruited and categorized into two groups based on abdominal obesity status: 22 individuals with



abdominal obesity and 14 individuals with a normal abdominal profile, with the flowchart available in [Figure 1](#). In this study, there were clear differences between people with abdominal obesity and those with non-abdominal obesity, particularly the abdominal obesity group had a higher BMI ($29.71 \pm 5.34 \text{ kg/m}^2$ vs. $23.32 \pm 4.20 \text{ kg/m}^2$), larger waist circumference ($98.83 \pm 11.11 \text{ cm}$ vs. $81.37 \pm 11.30 \text{ cm}$), and higher visceral fat ($12.32 \pm 3.44\%$ vs. $6.96 \pm 2.91\%$) compared to the normal obesity group. Full details are available in [Table 1](#).

3.2 Heatmap analysis

The heatmap analysis in [Figure 2](#) reveals distinct correlations between human health parameters and gut microbiota composition, with more than 80% represented by the following taxa: *Roseburia*

inulinivorans, *Faecalibacterium prausnitzii*, *Bacteroides massiliensis*, *Bacteroides caccae*, *Bacteroides stercoris*, *Prevotella stercorea*, *Bacteroides vulgatus*, *Phascolarctobacterium faecium*, *Collinsella aerofaciens*, *Bacteroides fragilis*, *Prevotella_9 copri*, *Sutterella wadsworthensis*, *Alistipes putredinis*, *Bacteroides uniformis*, *Parabacteroides merdae*, *Bacteroides plebeius*, and *Bacteroides coprocola*. In this analysis, significant correlations were observed between several species with biomarkers, with interpreted key details available in [Table 2](#).

3.3 Alpha and beta diversity

Alpha diversity analysis using species richness and diversity indices showed no significant differences between abdominal obesity




TABLE 1 Characteristics of individuals.

Subject characteristics	p-Value ^a	Abdominal obesity (n = 22)	Non-abdominal obesity (n = 24)
Age (years old)	0.360	35.27 ± 8.93	34.04 ± 9.68
Height (cm)	0.159	164.90 ± 10.72	160.71 ± 7.38
Weight (kg)	0.000*	81.47 ± 21.23	60.48 ± 12.66
BMI (kg/m ²)	0.000*	29.71 ± 5.34	23.32 ± 4.20
Waist circumference (cm)	0.000*	98.83 ± 11.11	81.37 ± 11.30
Fat mass (%)	0.005*	31.93 ± 6.57	25.94 ± 7.05
Muscle mass (%)	0.010*	27.62 ± 5.41	31.40 ± 5.05
Visceral fat (%)	0.000*	12.32 ± 3.44	6.96 ± 2.91
Systolic blood pressure (mmHg)	0.085	129.77 ± 16.56	121.67 ± 12.73
Diastolic blood pressure (mmHg)	0.097	89.18 ± 9.46	84.75 ± 8.71
Energy (kcal)	0.296	2043.02 ± 438.46	1907.65 ± 430.54
Protein (g)	0.257	88.76 ± 23.11	80.67 ± 22.14
Fat (g)	0.531	81.73 ± 21.00	77.30 ± 25.97
Carbohydrate (g)	0.350	245.89 ± 58.55	226.18 ± 52.86
Fiber (g)	0.982	17.45 ± 4.55	16.95 ± 5.17
PUFA (g)	0.296	20.56 ± 7.40	18.25 ± 6.09

^aMann–Whitney U Test.
All data are in Mean ± SD, with p-value of <0.05 (indicated with star (*)) being significant. BMI, Body mass index; PUFA, Polyunsaturated fatty acids.

TABLE 2 Species-to-biomarker key highlights from Spearman correlations analysis.

Variable	Species	Correlation value	p-Value ^a
BMI	<i>Prevotella_9 copri</i>	0.322	0.029
VF*		0.206	0.169
Diastolic blood pressure	<i>Bacteroides coprocola</i>	−0.302	0.032
PUFA		0.364	0.013
Energy	<i>Roseburia inulinivorans</i>	0.349	0.018
Protein		0.334	0.023
Fat	<i>Bacteroides massiliensis</i>	0.326	0.027
Carbohydrate	<i>Sutterella wadsworthensis</i>	0.294	0.047
Fiber	<i>Bacteroides plebeius</i>	0.388	0.008

^aAdjusted p-value.

*Indicating species-to-biomarker of interest. BMI, Body mass index; VF, Visceral fat; PUFA, Polyunsaturated fatty acid.

and non-abdominal obesity groups. Species numbers were similar ($p = 0.9$), and Shannon, Simpson, and Inverse Simpson indices showed no differences ($p = 0.72, 0.5, 0.5$), indicating similar microbial diversity between groups. Beta diversity analysis using NMDS and PCoA plots revealed no significant separation between groups, as supported by PERMANOVA results ($R^2 = 0.016, p = 0.953$), indicating minimal compositional differences in gut microbiota.

4 Discussion

This study identified several significant correlations among various species and biomarkers, as demonstrated through heatmap analysis in Figure 2 and detailed in Table 2. There are several potential metabolic functions of the identified microbes according to several studies. *Prevotella_9 copri* is notably involved in amino acid and carbohydrate metabolism (25), while *Bacteroides coprocola* has shown a characteristic distribution of single-nucleotide polymorphisms (SNPs) in type 2 diabetes patients (26). *Roseburia inulinivorans* is recognized for its utilization of inulin and starch (27), and *Bacteroides massiliensis* also plays a role in carbohydrate metabolism (28). *Sutterella wadsworthensis* demonstrates potential binding to mucus and extracellular matrix proteins (29), and *Bacteroides plebeius* has been isolated from seaweed-eating Japanese individuals (30). From these identified microbes, several correlations were observed. For instance, BMI exhibited a positive correlation with *Prevotella_9 copri*, consistent with findings from a study examining this relationship in children (31). While waist circumference correlated positively with *Prevotella_9 copri* and *Bacteroides coprocola*, no prior studies have reported similar findings. These species are abundant in individuals consuming carbohydrate-rich diets, suggesting a potential link between these species and waist circumference via carbohydrate intake. However, our study found no significant correlation between these species and carbohydrate consumption. (25, 32, 33). Several studies have reported findings that differ from those of the present study. For instance, Asnicar et al. (34) identified an inverse correlation between *Prevotella_9 copri* and visceral fat, whereas this study observed a positive correlation. This is interesting because while there is a possible explanation of increasing

PUFA serum level as the reason of negative correlation in one of the studies, in this study *Prevotella_9 copri* are having no significance with the PUFA serum level (34). Furthermore, there are few researches that explore the relationship between other species and biomarkers. This scarcity may be attributed to the limited exploration of species-biomarker correlation, especially in Indonesia, where dietary intake and other biomarkers differ from those in other regions.

Alpha diversity analysis showed no significant differences in species richness and diversity indices between groups, indicating maintained microbial biodiversity regardless of abdominal obesity status. Beta diversity analysis revealed no segregation of gut microbiota community structures between abdominal obesity and non-abdominal obesity groups, with NMDS and PCoA plots showing no clustering differences. These findings contrast with prior studies that demonstrated distinct clustering patterns between metabolically unhealthy individuals and healthy controls, suggesting obesity-linked microbial composition differences (35, 36). However, in this study, such clear distinctions were not observed. The lack of correlation between alpha and beta diversity may be explained by their different aspects: alpha diversity measures the richness and evenness within a single sample, whereas beta diversity measures the differences between samples. Study limitations include low sample size ($n = 46$) and cross-sectional design preventing causal inference. While taxonomic profiles were analyzed, microbial metabolite functions were not studied. Furthermore, other associated factors were not observed such as physical activity that showed the effects of the overall microbiota composition, limiting the generalizability (37). However, given the nature of this study as a pilot study, this work provides the groundwork for future microbiota research.

In conclusion, this study revealed a potential uniqueness in one of the gut microbiota, with *Prevotella_9 copri* showing an inverse correlation with visceral fat compared to findings from other studies conducted outside of Indonesia. These microbial profiles may serve as biomarkers for early detection and personalized interventions. However, the small sample size and cross-sectional design restrict the generalizability and causal inference. Future studies with larger sample sizes and comparisons across urban and rural locations, including coastal and highland regions, will be beneficial to validate these associations, enhance understanding of the underlying mechanisms, and advance microbiota-targeted strategies for managing abdominal obesity and its metabolic consequences.

Data availability statement

The original contributions presented in the study are included in the article/supplementary material; further inquiries can be directed to the corresponding author.

Ethics statement

The studies involving humans were approved by Medical Ethics Committee of the Faculty of Medicine, Diponegoro University. The studies were conducted in accordance with the local legislation and institutional requirements. The participants provided their written informed consent to participate in this study.

Author contributions

FF: Conceptualization, Methodology, Writing – original draft, Resources, Data curation, Writing – review & editing. FA: Visualization, Validation, Data curation, Formal analysis, Methodology, Writing – review & editing, Writing – original draft, Software. NM: Writing – review & editing, Data curation, Resources, Methodology, Formal analysis. JF: Writing – review & editing, Formal analysis, Methodology, Data curation, Resources. HZ: Methodology, Writing – review & editing, Formal analysis, Resources, Data curation. MS: Investigation, Methodology, Writing – review & editing, Resources, Data curation. HT: Conceptualization, Writing – review & editing. AW: Writing – review & editing, Validation. AJ: Validation, Writing – review & editing. AP: Writing – original draft, Funding acquisition, Resources, Supervision, Conceptualization, Investigation, Writing – review & editing.

Funding

The author(s) declare that financial support was received for the research and/or publication of this article. This study was partially funded by the Summit Institute for Development, the GSI Lab, and the Oxford University Clinical Research Unit Indonesia with support from the British Embassy in Jakarta as part of the Genomics and Science Dojo 2.0 program, for the Open Access publication fees.

Acknowledgments

We express our gratitude to Diponegoro University for their support in conducting this study through the World Class Research University (WCRU) grant program, administered by the Institute for Research and Community Services (LPPM), under WCRU program number 357-08/UN7. D2/PP/IV/2024. Furthermore, we acknowledge

the support and guidance provided by the Genomics and Science Dojo, and the Genomics and Science Workshop and facilitators during manuscript preparation. The Genomics and Science Dojo and Workshop was conducted by the Summit Institute for Development, the GSI Lab, and the Oxford University Clinical Research Unit–Jakarta with support from the British Embassy in Jakarta (FCDO Programme Name: Indonesia Digital Health, FCDO Project Number: 400422-409).

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Generative AI statement

The authors declare that no Gen AI was used in the creation of this manuscript.

Any alternative text (alt text) provided alongside figures in this article has been generated by Frontiers with the support of artificial intelligence and reasonable efforts have been made to ensure accuracy, including review by the authors wherever possible. If you identify any issues, please contact us.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Islam ANMS, Sultana H, Nazmul Hassan Refat M, Farhana Z, Abdulbasah Kamil A, Meshbahur RM. The global burden of overweight-obesity and its association with economic status, benefiting from STEPs survey of WHO member states: a meta-analysis. *Prev Med Rep.* (2024) 46:102882. doi: 10.1016/j.pmedr.2024.102882
- Ayuningtyas D, Kusuma D, Amir V, Tjandrarini DH, Andarwati P. Disparities in obesity rates among adults: analysis of 514 districts in Indonesia. *Nutrients.* (2022) 14:3332. doi: 10.3390/NU14163332
- Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the global burden of disease study 2013. *Lancet.* (2014) 384:766–81. doi: 10.1016/S0140-6736(14)60460-8
- Ferdina AR, Arfines PP, Aryastami NK. Obesity in urban Indonesia: evidence from the 2007 and 2018 basic health research. *Med J Indones.* (2024) 33:119–27. doi: 10.13181/MJI.OA.247183
- Indonesian Ministry of Health. Indonesian Health Survey in Number, Jakarta. (2023). Available online at: <https://www.badankebijakan.kemkes.go.id/en/ski-2023-dalam-angka/> (Accessed July 23, 2025)
- Geiger SJ, White MP, Davison SMC, Zhang L, McMeel O, Kellett P, et al. Coastal proximity and visits are associated with better health but may not buffer health inequalities. *Commun Earth Environ.* (2023) 4:166. doi: 10.1038/s43247-023-00818-1
- Fan Y, Pedersen O. Gut microbiota in human metabolic health and disease. *Nat Rev Microbiol.* (2020) 19:55–71. doi: 10.1038/s41579-020-0433-9
- Karwur FF, Yocku MHSO, Enoch DA, Triandhini RLNR, Agustina V, Lakukua MF, et al. Anthropometric and metabolic differences and distribution of ABCG2 rs2231142 variant between lowland and highland Papuans in West Papua, Indonesia. *J Physiol Anthropol.* (2025) 44:14. doi: 10.1186/s40101-025-00394-7
- Nurhasan M, Ariesta DL, Utami MMH, Fahim M, Aprillyana N, Maulana AM, et al. Dietary transitions in Indonesia: the case of urban, rural, and forested areas. *Food Secur.* (2024) 16:1313–31. doi: 10.1007/s12571-024-01488-3
- Kootte RS, Levin E, Salojärvi J, Smits LP, Hartstra AV, Udayappan SD, et al. Improvement of insulin sensitivity after lean donor feces in metabolic syndrome is driven by baseline intestinal microbiota composition. *Cell Metab.* (2017) 26:611–619.e6. doi: 10.1016/j.cmet.2017.09.008
- Zhao L, Zhang F, Ding X, Wu G, Lam YY, Wang X, et al. Gut bacteria selectively promoted by dietary fibers alleviate type 2 diabetes. *Science.* (2018) 359:1151–6. doi: 10.1126/SCIENCE.AAO5774
- Fiçıcılar BB. Comparative analysis of fish consumption habits in coastal and inland districts of Samsun Province. *Mar Sci Technol Bull.* (2024) 13:251–61. doi: 10.33714/MASTEB.1524232
- Bratlie M, Hagen IV, Helland A, Erchinger F, Middtun Ø, Ueland PM, et al. Effects of high intake of cod or salmon on gut microbiota profile, faecal output and serum concentrations of lipids and bile acids in overweight adults: a randomised clinical trial. *Eur J Nutr.* (2020) 60:2231–48. doi: 10.1007/S00394-020-02417-8
- Casari S, Di Paola M, Banci E, Diallo S, Scarallo L, Renzo S, et al. Changing dietary habits: the impact of urbanization and rising socio-economic status in families from Burkina Faso in sub-Saharan Africa. *Nutrients.* (2022) 14:1782. doi: 10.3390/NU14091782/S1
- Tham KW, Abdul Ghani R, Cua SC, Deerochanawong C, Fojas M, Hocking S, et al. Obesity in south and Southeast Asia—a new consensus on care and management. *Obes Rev.* (2023) 24:e13520. doi: 10.1111/obr.13520
- World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA.* (2013) 310:2191–4. doi: 10.1001/JAMA.2013.281053

17. Ross R, Neeland IJ, Yamashita S, Shai I, Seidell J, Magni P, et al. Waist circumference as a vital sign in clinical practice: a consensus statement from the IAS and ICCR working group on visceral obesity. *Nat Rev Endocrinol.* (2020) 16:177–89. doi: 10.1038/s41574-019-0310-7
18. Ramírez-Vélez R, Tordecilla-Sanders A, Correa-Bautista JE, González-Ruiz K, González-Jiménez E, Triana-Reina HR, et al. Validation of multi-frequency bioelectrical impedance analysis versus dual-energy X-ray absorptiometry to measure body fat percentage in overweight/obese Colombian adults. *Am J Hum Biol.* (2018) 30:1–6. doi: 10.1002/AJHB.23071
19. Syaury A, Afifah DN, Purwanti R, Nissa C, Fitranti DY, Chao JCJ. Reproducibility and validity of a food frequency questionnaire (FFQ) developed for middle-aged and older adults in Semarang, Indonesia. *Nutrients.* (2021) 13:4163. doi: 10.3390/NU13114163
20. Herforth AW, Ballard T, Rzepa A. Development of the diet quality questionnaire for measurement of dietary diversity and other diet quality indicators. *Curr Dev Nutr.* (2024) 8:103798. doi: 10.1016/j.cdnut.2024.103798
21. Lim MY, Park Y-S, Kim J-H, Nam Y-D. Evaluation of fecal DNA extraction protocols for human gut microbiome studies. *BMC Microbiol.* (2020) 20:212. doi: 10.1186/s12866-020-01894-5
22. Nearing JT, Douglas GM, Hayes MG, Mac Donald J, Desai DK, Allward N, et al. Microbiome differential abundance methods produce different results across 38 datasets. *Nat Commun.* (2022) 13:1–16. doi: 10.1038/s41467-022-28034-z
23. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol.* (1995) 57:289–300. doi: 10.1111/J.2517-6161.1995.TB02031.X
24. Li R-J, Jie Z-Y, Feng Q, Fang R-L, Li F, Gao Y, et al. Network of interactions between gut microbiome, host biomarkers, and urine metabolome in carotid atherosclerosis. *Front Cell Infect Microbiol.* (2021) 11:708088. doi: 10.3389/fcimb.2021.708088
25. Abdelsalam NA, Hegazy SM, Aziz RK. The curious case of *Prevotella copri*. *Gut Microbes.* (2023) 15:2249152. doi: 10.1080/19490976.2023.2249152
26. Chen Y, Li Z, Hu S, Zhang J, Wu J, Shao N, et al. Gut metagenomes of type 2 diabetic patients have characteristic single-nucleotide polymorphism distribution in *Bacteroides coprocola*. *Microbiome.* (2017) 5:15. doi: 10.1186/s40168-017-0232-3
27. Scott KP, Martin JC, Chassard C, Clerget M, Potrykus J, Campbell G, et al. Substrate-driven gene expression in *Roseburia inulinivorans*: importance of inducible enzymes in the utilization of inulin and starch. *Proc Natl Acad Sci.* (2011) 108:4672–9. doi: 10.1073/pnas.1000091107
28. Pudlo NA, Urs K, Crawford R, Pirani A, Atherly T, Jimenez R, et al. Phenotypic and genomic diversification in complex carbohydrate-degrading human gut bacteria. *mSystems.* (2022) 7:e0094721. doi: 10.1128/msystems.00947-21
29. Davoody S, Halimi H, Zali A, Houri H, Brand S. Double-edged sword effect of Sutterella in neurological disorders: implications for the gut-brain axis and neuroimmune interactions. *Neurobiol Dis.* (2025) 214:107032. doi: 10.1016/J.NBD.2025.107032
30. Park NJ, Yu S, Kim DH, Yun EJ, Kim KH. Characterization of BpGH16A of *Bacteroides plebeius*, a key enzyme initiating the depolymerization of agarose in the human gut. *Appl Microbiol Biotechnol.* (2021) 105:617–25. doi: 10.1007/s00253-020-11039-3
31. Gong J, Zhang Q, Hu R, Yang X, Fang C, Yao L, et al. Effects of *Prevotella copri* on insulin, gut microbiota and bile acids. *Gut Microbes.* (2024) 16:1–12. doi: 10.1080/19490976.2024.2340487
32. Lewis S, Nash A, Li Q, Ahn TH. Comparison of 16S and whole genome dog microbiomes using machine learning. *BioData Min.* (2021) 14:41. doi: 10.1186/s13040-021-00270-X
33. Cho YA, Choi JH. Association between carbohydrate intake and the prevalence of metabolic syndrome in Korean women. *Nutrients.* (2021) 13:3098. doi: 10.3390/NU13093098/S1
34. Asnicar F, Berry SE, Valdes AM, Nguyen LH, Piccinno G, Drew DA, et al. Microbiome connections with host metabolism and habitual diet from 1,098 deeply phenotyped individuals. *Nat Med.* (2021) 27:321–32. doi: 10.1038/s41591-020-01183-8
35. Le Chatelier E, Nielsen T, Qin J, Prifti E, Hildebrand F, Falony G, et al. Richness of human gut microbiome correlates with metabolic markers. *Nature.* (2013) 500:541–6. doi: 10.1038/nature12506
36. Geng J, Ni Q, Sun W, Li L, Feng X. The links between gut microbiota and obesity and obesity related diseases. *Biomed Pharmacother.* (2022) 147:112678. doi: 10.1016/J.BIOPHA.2022.112678
37. Monda V, Villano I, Messina A, Valenzano A, Esposito T, Moscatelli F, et al. Exercise modifies the gut microbiota with positive health effects. *Oxidative Med Cell Longev.* (2017) 2017:3831972. doi: 10.1155/2017/3831972